



International Conference
on
**NATURAL SCIENCE AND GREEN TECHNOLOGIES
FOR SUSTAINABLE DEVELOPMENT (NTSD-2022)**

30th November, 1st and 2nd December, 2022
Goa University, Taleigao Plateau, Goa

Jointly Organized by



National Environmental Science Academy
New Delhi

School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Abstracts and Souvenir

Sponsored by





Please visit us at
www.nesa-india.org




International Conference
on
**NATURAL SCIENCE AND GREEN TECHNOLOGIES
FOR SUSTAINABLE DEVELOPMENT (NTSD-2022)**

30th November, 1st and 2nd December, 2022
Goa University, Taleigao Plateau, Goa

Jointly Organized by



National Environmental Science Academy
New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Abstracts and Souvenir

Sponsored by



Message



I am pleased to learn that the School of Biological Sciences and Biotechnology, Goa University and National Environmental Science Academy (NESA), New Delhi is organising an international conference on "Natural Science and Green Technologies for Sustainable Development (NTSD-2022)" from 30th November to 2nd December 2022.

I am glad to note that this conference will create an ecosystem for eminent scholars to discuss, interact and give impetus to research and extension activities. Key issues will be discussed, which are directly associated with the current trend in green technology dealing with challenges and prospects of society, where living conditions and resources are used to continue to meet human needs without undermining the integrity and stability of the natural system.

The objective of this conference is to focus on technological advancement to curb the environmental pollution and propose solutions to adopt the green technologies for sustainable development.

I am sure, this conference will provide great opportunities to the participants and serve as an effective platform for exchanging ideas.

I congratulate the Organisers and wish the Conference a grand success. I wish the participants a very enriching and enlightening experience and my greetings to all the renowned speakers of this Conference.


Harilal B. Menon 15/11/2022
Vice-Chancellor and
Patron- NTSD 2022



National Environmental Science Academy

(Registered Under Society Act XXI of 1860)

206, Raj Tower-I, Alaknanda Community Centre, New Delhi-110 019

Phone: 011-2602 3614 (O)

E-mail: infonesa88@gmail.com; nesapublications@gmail.com

22.11.2022

Message



5I and the members of the organizing committee are very proud to present the International Conference on NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR SUSTAINABLE DEVELOPMENT (NTSD-2022) welcoming all the participants to Goa University, Goa from 30th November to 2nd December 2022.

It gives me immense pleasure to welcome you all as President of NESA.

The present conference that is being jointly organized by NESA, New Delhi and Goa University, Goa.

I also heartily welcome and congratulate all the Awardees of NESA – 2022 who will receive their respective awards on 30th November and 2nd December, 2022.

The conference is aimed at expanding the programme by including all aspects related to safe agriculture, environment, forest, ecosystems, food, climate change, sustainability of resources, renewable and non-renewable energy, global warming, extinction of species, malnutrition, hunger, drought & famine, biotechnological tools, integrating components for production and protection as a multidisciplinary approach for essential sustainable management.

The various sub-themes of the conference will offer many opportunities to delegates to learn many things and apply the same in their respective workplace.

I am sure the young scientists will get benefit after listening the invited talks in this conference.

I would also like to put before you the related recommendations of the renowned COP27 Climate Summit - November 6 – 18, 2022 at SHARM EL-SHEIKH, Egypt. For the protection of environment 50 billion trees will be planted under the Green Initiative in Africa as per joint declaration in the above summit.

Protection of oceans, clean energy, food availability, reduction of greenhouse gasses, carbon and fossil fuel are the recommendations of COP27. Thus the global warming and the increase of temperature of 1.5 degrees Celsius may be checked upto 2030. As per U.N. last eight years remained hot in the history of the world. Protection of endangered species & ecosystems (Congo forest, Gorilla population, Mangroves forest and coral reefs) is declared in joint declaration of COP27. Judicious use of resources is the key in climate fight.

Please visit our website : www.nesa-india.org

The world population has become 8 billion in November, 2022 as per annual report issued by United Nations. It will be a great platform to strengthen the friendship and collaboration among scientists, academics and industries.

The diversity of specializations and current themes and issues, will enable us to achieve our targeted mandate and vision. Thus India may be able to achieve key goals to fight against climate change, food shortage and less emissions of Methane and GHGs.

We will have awardees, guests, keynote speakers, from the different states including Goa from varied fields of environment, agriculture, ecosystem, food, energy and Biotechnology / Biosciences.

The hard work and dedication of all the members of organizing, scientific, technological, hospitality, transport, boarding, lodging and financial committees during the preparation of this conference is highly appreciated at may end.

I am grateful to Prof. Harilal B. Menon, Vice Chancellor, Goa University for giving permission to hold the conference here as well as his Patronage for the success of conference.

Prof. (Dr.) Savita Kerkar, Dean, School of Biological Sciences & Biotechnology, Goa University, Goa and Convener NTSD 2022 is highly thanked for her valuable support and supervision on all activities of the conference.

Dr. Mrutyunjay Suar is also thanked for his continued encouragement during organizing the present event.

Prof. (Dr.) Shakeel Ahmad Khan, Organizing Secretary is also highly appreciated for his keen interest and continued efforts for the success of the conference.

Financial assistance from the Govt. agencies and N.G.O.'s for the present conference is also highly acknowledged.

The staff of NESAMr. Gian Chand and Mr. Rakesh Kumar Roy also deserve appreciation for their hard work and sincerity.

Without them, the event would not have been possible.

My personal respect goes out to all of you.

I look forward for the success of NTSD 2022.



(Prof. Javed Ahmd)



गोंय विद्यापीठ

ताळगांव पठार,

गोंय - ४०३ २०६

फोन : +९१-८६६९६०९०४८

फॅक्स : +०९१-८३२-२४५९९८४/२४५२८८९



Goa University

Taleigao Plateau, Goa-403 206

Tel : +91-8669609048

Fax : +091-832-2451184/2452889

E-mail : registrar@unigoa.ac.in

Website : www.unigoa.ac.in

(Accredited by NAAC with Grade 'A')

Message



Natural Sciences decipher the rules that govern the natural world through scientific methods. Since ancient times, natural sciences has been changing our lives and ways of thinking at a much greater pace than in the earlier period of human civilization. A natural science, viz Green Technology or clean technology, deals with applying environmental science, green chemistry, environmental monitoring and sensors to monitor, model and conserve the existing natural environment and its resources. The history of natural sciences is inseparable from the gradually increasing awareness and understanding of indirect effects. Indirect interactions are not apparent, and their studies often face challenges of system complexity. However, the knowledge of complex interactions involving ecosystem components is indispensable for the sustainable development of humankind. Systematic elucidation of the indirect effects is becoming a central focus for ecology and environmental sciences.

Thus, the International Conference on Natural Science and Green Technologies for Sustainable Development (NTSD-2022) is a timely endeavour to carve out a futuristic, economically viable, and environmentally friendly solution. Being the contributor to organizing this conference, I anticipate a declaration at the end of the meeting that could be prescribed to the environmental agency. The conference will provide a unique platform to learn about the advancement in green technology and initiatives that are being taken to maintain and improve the quality of the environment and give new resource-efficient and sustainable thinking to the society.

Savita Kerkar
(Savita Kerkar)



National Environmental Science Academy

(Registered Under Society Act XXI of 1860)

206, Raj Tower-I, Alaknanda Community Centre, New Delhi-110 019

Phone: 011-2602 3614 (O)

E-mail: infonesa88@gmail.com; nesapublications@gmail.com

21.11.2022

Message



Dear delegates and friends Warm greetings!!!

On behalf of NESA and the organizing committee, I would like to cordially welcome you to the International conference on "Natural Science and Green Technologies for Sustainable Development" to be held at Goa University, Goa from the 30th November to 2nd Dec., 2022.

We had been working for the natural resources management and sustainable development for the last 5 decades to create awareness and educate people to realise that sustainable development is the keys factor linking to the quality of life of people and their ecosystem. We could successfully associate ourselves with various technology institutions and academia in the last decades and successfully demonstrated that environmental management and sustainable development could be achievable by joining hands together. This conference with the theme "Natural Science and Green Technologies for Sustainable Development" is the right platform to bring various academicians under one roof to discuss needs of current scenario to mitigate the global warming and pollution problem by interlink the sustainable development goals of UNEP.

We are planning to have best interaction with multi domain displays and poster presentation. The thematic talks and the plenary sessions will drive you through the multi sectoral emergence and mitigation options to achieve SDGs. This could be the first conference of its kind in the region where everyone could have opportunity to showcase and present their ideas, thoughts, developments that could lead to a meaningful life in the community. We are trying our best to ensure that your time and stay in Goa during the conference be one of the most memorable one and you go back with rich information.

I welcome you, your family and friends again to this wonderful gathering and make the maximum out of it. I thank each and every one of you who are contributing to the success of the conference and looking forward to seeing you all soon.

JAI HIND

Best wishes

(Dr. Shakeel Ahmad Khan)

General Secretary NESA

Principal Scientist, Division of Environmental Sciences, ICAR-IARI, New Delhi -110012

Organising Secretary – NTSD 2022

Please visit our website : www.nesa-india.org

About the Organizers

The School of Biological Sciences and Biotechnology (SBSB), Goa University, Goa

The School of Biological Sciences and Bio-technology (SBSB) was established in May, 2022 merging four Disciplines to form the School of Biological Sciences and Biotechnology. Goa being a coastal state, in proximity to the Western Ghats, a biodiversity hot-spot, provides several unique ecosystems for research.

The focus of amalgamation of the four departments was to bring fundamental and biological science Departments under one umbrella to provide the students a platform for interdisciplinary research.

Individually in the past, the departments have evolved in a step wise manner. The Department of Microbiology established in 1974 was a part of the Centre of Post-graduate Instructions and Research (CPIR) of the University of Mumbai in 1985, it became a full-fledged Department of the then newly founded Goa University. The Department of Botany and the Department of Zoology, were both established in 1990, became full-fledged Departments of the Goa University providing students with an opening for the Master's programme in Botany and Zoology. The Department of Biotechnology, one of the pioneer in Marine Biotechnology (1998) teaching and research in the country, became full-fledged Department of the Goa University in the year 2003. Subsequently, a new M.Sc. and Ph.D. Programme in Biotechnology were established. The Marine Biotechnology programme has been continuously funded by the DBT, Government of India. The four departments in the School of SBSB have come together to multiply their strengths and carry out effective interdisciplinary teaching and research in biological sciences. Equally, an important aspect of biological sciences, namely Home Sciences and Wildlife studies have been integrated to further gain in-depth knowledge of biological sciences.

The focus of the school is to provide quality education and research in biological sciences through an integrated approach with a view to create well trained human resource to handle the upcoming challenges in a sustainable manner and to apply the knowledge to the welfare of the society.

National Environmental Science Academy (NESA), New Delhi

National Environmental Science Academy (NESA) was founded by Late Prof. TRC Sinha, the then Head of Zoology Department, MJK PG College, Bihar University to create awareness, promote and protect the environment. Conceptualised and initiated in 1984, it was registered as a Society in 1988 under the Societies Act XXI of 1860 at Patna.

This ACADEMY is of National level, presently having its Head Office at 206, Raj Tower-1, Alaknanda Community Centre, New Delhi. The main objective of the Academy is to bring awareness about the environmental issues among the masses and strive to find sustainable solutions by arranging lectures, demonstrations, training programmes, seminars, symposium, conferences, publishing journals and organizing any other activities supporting the cause.

Aims Objectives and Functions of the Academy

To implements the SDDGs in India by encouraging students, scientists, researchers, academicians and members of the academy for pursuing research on sustainable development.

- **To set up Regional/State Chapters for dissemination of information on environment.**
- **To motivate and prepare young minds on environmental management.**
- **To organize national/international level conferences, symposia, seminars, meetings and workshops on themes of environmental concerns.**
- **To publish policy papers, synthesis volumes, proceedings, journals, newsletter, transactions and other publications for the promotion of Environmental Sciences.**
- **To forward the recommendation of scientists /professors to govt. agencies.**

Various eminent personalities have graced the Academy as President. The first President of the Academy was Dr. K.C. Bose, Vice-Chancellor of Ranchi University; then Dr. B.S. Attri, Advisor, Ministry of Environment and Forest. Most recently Padma Bhushan Dr. S.Z. Qasim was the President of the Academy till June 2015, who is a renowned marine scientist known for his Antarctica mission in 1981-82, he also served as the Secretary at the Dept. Of Ocean Development (now Ministry of Earth Sciences); Member, Planning Commission and Vice-Chancellor, Jamia Millia Islamia, New Delhi. Currently Prof. Javed Ahmad, (Former Dean, Faculty of Science), Jamia Hamdard, New Delhi, is the President of the Academy.

Annual Awards

The Academy recognizes the merit and achievements of individuals who have contributed to the field of environmental science, education and societal values by conferring **(1) NESA FELLOWSHIP OF THE YEAR AWARD (2) NESA EMINENT SCIENTIST OF THE YEAR AWARD (3) NESA SCIENTIST OF THE YEAR AWARD (4) NESA ENVIRONMENTALIST OF THE YEAR AWARD (5) NESA GREEN TECHNOLOGY INNOVATIVE AWARD (6) NESA DISTINGUISHED SCIENTIST OF THE YEAR AWARD (7) WOMEN EXCELLENCE OF THE YEAR AWARD (8) NESA YOUNG SCIENTIST OF THE YEAR AWARD (9) NESA JUNIOR SCIENTIST OF THE YEAR AWARD (10) NESA BEST RESEARCHER AWARD (11) NESA BEST PHD THESIS AWARD.**

Any life member of the Academy can apply for the awards. For more information please, log on to our website: <http://nesa-india.org/award-form-submission>.

In addition, the Best Oral Presentation and Best Poster Awards are given away during the Annual Conference of the Academy.

Publications

The Academy is publishing the following Journals (Biannual):

- 1) **INTERNATIONAL JOURNAL ON AGRICULTURAL SCIENCES**
- 2) **INTERNATIONAL JOURNAL ON ENVIRONMENTAL SCIENCES**
- 3) **INTERNATIONAL JOURNAL ON BIOLOGICAL SCIENCES**
- 4) **INDIAN JOURNAL OF UNANI MEDICINE**
<http://nesa-india.org/nesa-journal/>
- 5) **E-NESA Newsletter (Monthly)** <http://nesa-india.org/newsletter/>

PATRON'S

Prof. Harilal B. Menon, Vice Chancellor, Goa University, Taleigao Plateau Goa

Prof. Javed Ahmad, President, National Environmental Science Academy, Delhi

CO-PATRONS

Dr. R.C. Agrawal, Deputy Director General (Agricultural Education), ICAR, New Delhi

Dr. R. Basavarajappa, Vice Chancellor, University of Agricultural Sciences, Dharwad

CONVENER

Prof. Savita Kerkar, Dean, School of Biological Sciences and Biotechnology, Goa University, Taleigao Plateau, Goa, India

ORGANIZING SECRETARY

Dr. Shakeel Ahmad Khan, Professor/Principal Scientist, Division of Environmental Sciences, NRL Building, Pusa Campus ICAR- Indian Agricultural Research Institute, New Delhi, India

SCIENTIFIC COMMITTEE

Dr. P. S. Brahmanand, Director of Research, RPCAU, Pusa, Samastipur, Bihar

Dr. Tanu Jindal, Group Additional Pro Vice Chancellor (R&D), J1 , G-21, Amity University, Sector 125, Noida, UP, India

Dr. S. S. Rathore, Principal Scientist, Division of Agronomy & Coordinator-NAHEP, IARI, Pusa, New Delhi

Dr. A.K. Pandey, Ex Principal Scientist NBFGR, Lucknow

Prof. Boyina Rupini, SOITS, Indira Gandhi National Open University, New Delhi

Dr. S V Sai Prasad, Principal Scientist, Department of Plant Breeding, ICAR-Indian Institute of Rice Research, Rajendranagar, Hyderabad.

Prof. Debidas Ghosh, Head Department of Bio-Medical Laboratory Science & Management Vidyasagar University, West-Bengal

Dr. Manoj Shrivastava, Principal Scientist, Division of Environment Science, NRL, Indian Agricultural Research Institute, Pusa, New Delhi

Dr. Pankaj Kumar Srivastava, Principal Scientist, Environmental Technologies Division, CSIR-National Botanical Research Institute, Lucknow, U.P.

Dr. Raaj K. Maheshwari, Former Professor Department of Chemistry SBRM Govt PG College, Nagaur, Rajasthan & Freelance Investigator & Scientific Writer
Dr. Mukesh Kumar Gupta, National Institute of Technology Rourkela, Odisha
Dr. M. Suresh Gandhi, Dept. of Geol., Head, Center for Env. Sciences, University of Madras, Chennai
Dr. Amal Chandra Mondal, Associate Professor, School of Life Sciences, Jawaharlal Nehru University, Delhi
Dr. Suman Mor, Associate Professor, Department of Environment Studies, Panjab Univeristy, Chandigarh
Dr. Rishi Saxena, Associate Professor, Department of Microbiology Bundelkhand University Jhansi, U.P.
Dr. Richa Rai, Senior Scientist, CSIR-NATIONALBOTANICALRESEARCH INSTITUTE, Lucknow, U.P.
Dr. Asif Iquebal, Senior Scientist, ICAR-Indian Agricultural Statistics Research Institute, New Delhi
Dr. Sarika, Senior Scientist, ICAR-Indian Agricultural Statistics Research Institute, New Delhi
Dr. V Sreeharsha Rachapudi, Asst. Prof., Dept. of Life Sc. and Biotechnology, CSJM University, Kanpu, U.P.
Dr. Jitendra K. Nagar, Environment & Social Development Association (ESDA) Delhi & Ass. Prof., Dr Bhim Rao Ambedkar College, University of Delhi, Delhi
Dr. Pradeep Kumar, Department of Zoology, Govt. Degree College Skukrauli, Kushinagar (U.P.)
Dr. Deeksha Dave, Asst. Prof., Dept. of Environmental, SOITES Disciplinary Studies, IGNOU, New Delhi
Dr. Surender S. Yadav, Asst. Prof., Dept. of Botany, Maharshi Dayanand University, Rohtak (Haryana)
Dr. J. Sangeetha, Assistant Professor, Dept. of Environmental Science, Central University of Kerala, Kerala
Dr. Lalit Mohan, Assistant Professor Department of Zoology, Dayalbagh Educational Institute (Deemed to be University), Agra, Uttar Pradesh
Dr. Sankha Chakraborty, Assistant Professor, Kalinga School of Biotechnology/Chemical Technology Kalinga Institute of Industrial Technology (Deemed to be University), Bhubaneswar, Odisha
Dr. S. Ezil Vedan, Scientist, CSIR-Central Food Technological Research Institute, Mysore, Karnataka
Dr. Asit Mandal, Sr. Scientist, ICAR-Indian Institute of Soil Science, Bhopal, Madhya Pradesh
Dr. Bharath Gandu, Assistant Professor, Department of Environmental Studies, University of Delhi, Delhi
Dr. Chandrakumar Kalichamy, Dept. of Renewable Energy, TNAU, Lawley, Coimbatore
Dr. Sushil Kumar Kharia, Department of Soil Science, College of Agriculture, SKRAU, Bikaner.

ORGANIZING COMMITTEE

Prof. Sanjeev C. Ghadi, Vice-Dean (Academics), School of Biological Sciences and Biotechnology, Goa University, Goa.
Prof. S. Krishnan, Vice-Dean (Research), School of Biological Sciences and Biotechnology, Goa University.
Dr. Nitin Sawant, School of Biological Sciences and Biotechnology, Goa University, Goa.
Dr. Milind M. Naik, School of Biological Sciences and Biotechnology, Goa University, Goa.
Dr. Shanti N. Desai, School of Biological Sciences and Biotechnology, Goa University, Goa.
Dr. Vikash Kumar Yadav, School of Biological Sciences and Biotechnology, Goa University, Goa.
Dr. Bhakti Balkrishna Salgaonkar, School of Biological Sciences and Biotechnology, Goa University, Goa.
Dr. Siddhi K. Jalmi, School of Biological Sciences and Biotechnology, Goa University, Goa
Ms. Dviti D. Mapari, School of Biological Sciences and Biotechnology, Goa University, Goa
Ms. Vinita, Division of Environmental Sciences, NRL Building, Pusa Campus ICAR-IARI, Delhi
Ms. Lukeshwari Shyam, Div. of Environmental Sciences, NRL Building, Pusa Campus ICAR-IARI, Delhi

NESA ORGANIZING COMMITTEE

Prof. A. Wanganeo, Vice President, Delhi & Former, Head DES & Limn., Director, UIT, B. University, Bhopal.
Prof. Altaf Ahmad, Vice President, NESA & Dep. of Botany, AMU, Aligarh, U.P.
Dr. Mridul Sahani, Vice President, NESA & Dr. B Sahni Clinic, Shivpuri Patna, Bihar
Dr. Prabhakar Ranjan, Joint Secretary, NESA, Delhi
Mr. RK Sinha, Executive Secretary, NESA, Delhi & Chairman Prajna Foundation, Delhi
Dr. Shri Prakash, Mem. Ex. Comm., NESA & Dept. of Zoology, K.A.P.G. College Allahabad
Dr. Ashok Dhakad, Mem. Ex. Comm., NESA & Scientist (Forestry), Chamber No. 131-B, Department of Forestry & Natural Resources, Punjab Agricultural University, Ludhiana, Punjab,
Dr. Balwant Rawat, Mem. Ex. Comm., NESADelhi & Asst. Prof., Dept. of Agriculture and Forestry, Graphic Era Hill University, Dehradun, Uttarakhand, India
Dr. Sandeep Kumar, Scientist, Department of Environmental Sciences, E-43, Pusa Campus, IARI, New Delhi
Mrs. Vandana Sinha, Mem. Executive Committee, NESADelhi & Social Worker, Delhi
Dr. Mamata Kumari, Ex. Comm., NESA & R.D.S. College, Muzaffarpur
Dr. Suraj K. Tripathy, Ex. Comm., NESA & Associate Dean (Academics), School of Chemical Technology, Associate Professor, School of Biotechnology, KiiT, Bhubaneswar.





Abstracts

International Conference on **NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR SUSTAINABLE DEVELOPMENT (NTSD-2022)**

30th November, 1st and 2nd December, 2022
Goa University, Taleigao Plateau, Goa

Contents

Sl. No.	Title & Author of the Abstract	Page No.
Technical Session 1		
1.	BIOMONITORING OF INDOOR AIR CONTAMINANTS USING SPIDER WEBS Aradhana K S., M. Vishnu Sreejith and Usha K Aravind	3
2.	IMPACT OF ENVIRONMENT AND LIFE STYLE ON GYNAECOLOGICAL CANCER RISK Nirmala K. and Manjulatha C.	4
3.	ASSESSMENT OF GROUNDWATER CONTAMINATION IN COASTAL AQUIFERS IN NORTH GOA, GOA, INDIA Mahesh Mayekar, Trupti Asolkar and A. Viegas	5
4.	BASELINE PATTERN OF MICROPLASTICS, THE OMNIPRESENT POLLUTANT IN THE WATERS OF VEMBANAD KOL WETLAND OF KERALA, INDIA Nisari A R and Sujatha C H	6
5.	CONSTITUENTS OF MAJOR XENOBIOTICS IN THE FISHERY HUB, VYPIN, KOCHI Jayakrishnan Unnikrishnan, Lisa Jose, Emil John C.R., Jyoti Varier, Nisari A.R., and Sujatha C.H.	7
6.	CONTINUOUS MAPPING OF RIVERINE SYSTEM IN KOCHI CITY USING MASS SPECTROMETRY Amitha Unnikrishnan, Usha K Aravind, Shiny Thomas and C. T. Aravindakumar	8
7.	DISTRIBUTION PROFILE OF BIOCHEMICAL CONSTITUENTS IN THE CORE SEDIMENTS OF VYPIN, THE FISHING HARBOUR OF KOCHI Emil John C.R., Jayakrishnan Unnikrishnan, Lisa, Nisari A.R., Jyoti Varrrior and Sujatha C H	9
8.	ESSENTIAL MINERAL AND HEAVY METALS AVAILABILITY IN VARIOUS WATER RESOURCES OF COLD DESERT HIGH-ALTITUDE REGION OF LADAKH, INDIA Vijay K. Bharti, Guru Charan, Arup Giri, Sudhir Kumar and OP Chaurasia	10
9.	PREVALENCE AND ASSESSMENT OF FLUORIDE LEVELS IN GROUNDWATER OF HINGOLI, MAHARASHTRA Pushyamitra Joshi, N. G. Shinde and P. P. Joshi	11
10.	ADDRESSING EARLY BLOOMING IN ALMONDS UNDER NORTH-WESTERN HIMALAYAN REGION OF KASHMIR Susheel Kumar Raina, Javed Iqbal Mir, Om Chand Sharma and Sheikh M Sultan	12
11.	ANTI-DIABETIC PROPERTIES OF MARINE OLIGOSACCHARIDES Devika Jadhav and Sanjeev C. Ghadi	13
12.	CHARACTERIZATION OF SEAWEED POLYSACCHARIDE DEGRADING BACTERIA AND THEIR EXPLOITATION FOR ANTI-DIABETIC ACTIVITY Elaine Antoinette Da Costa	14
13.	ECO-FRIENDLY APPROACH FOR DETOXIFICATION OF A HARMFUL TEXTILE DYE AND ITS PHYTOTOXIC EVALUATION Priyanka Bhagwan Jadhavar	15

Sl. No.	Title & Author of the Abstract	Page No.
14.	ELEVATED CO2 AFFECTS DNA METHYLATION PATTERN OF LOW AND HIGH ELEVATION POPULATIONS OF ARABIDOPSIS THALIANA Ashwani Kumar Verma, Akanksha Singh, Rajneesh Singh, Prabhatam Mishra, Shiv Narayan, Vivek Pandey, Pramod Arvind Shirke and Sribash Roy	16
15.	ELUCIDATION OF MICROBIAL SIDEROPHORES IN SALT TOLERANT KORGUT RICE VARIETY OF GOA Savia Bernadette Botelho	17
16.	ENZYMATIC HYDROLYSIS OF CASTOR OIL BY A NEWLY ISOLATED MICROBIAL STRAIN Shikha Singh, Sumit Sharma, Saurabh Jyoti Sarma and Satinder Kaur Brar	18
17.	EVALUATION OF METAL TOLERANCE OF HALOPHILIC EUBACTERIAL AND ARCHAEAL STRAINS ISOLATED FROM SOLAR SALTERNS OF GOA INDIA Fazila Shaikh and Sanika Naik-Samant	19
18.	INTEGRATED FABRICATION OF A BIO-BASED COMPOSITE BY LACTIC ACID CONDENSATION WITH RICE STRAW/LIGNIN FOR MAKING SUSTAINABLE GOODS AND UTILITY ITEMS Sumit Sharma, Shikha Singh and Saurabh Jyoti Sarma	20
19.	METAL-ORGANICFRAMEWORK BASED ADSORBENTS FOR THE REMOVAL OF DICLOFENAC FROM SYNTHETIC WATER Amrutha Asokana and C.T. Aravindakumar	21
20.	MICROPLASTIC TOXICITY: AN ANALYTICAL INSIGHT INTO DETECTION METHODS AND MYCO-DEGRADATION PATHWAYS Swati Solanki and Rachana Singh	22
21.	PHYSIOLOGICAL RESPONSE OF ECTOMYCORRHIZAL FUNGUS PISOLITHUS SP. ISOLATED FROM MINING SITE TO TEMPERATURE STRESS Manasi Kamat and Lakshangy Charya	23
22.	UTILIZATION OF AGRO-RESIDUE IN HIGH-PERFORMANCE SUPERCAPACITORS: A BRIEF CRITICAL REVIEW Ravi Kumar Sahu, S. Gangil and V.K. Bhargav	24
23.	CARABID BEETLES (COLEOPTERA: CARABIDAE) IN THE RICE AGRO-ECOSYSTEM OF KERALA Ashly Kurian and Sabu K. Thomas	25
24.	EDIBLE FILM PACKAGING-A NOVEL APPROACH IN FOOD PACKAGING Abhipsa Khobragade and Sumiran Dabrase	26
25.	EFFECT OF NATURAL ANTI-BROWNING AGENTS IN AVOIDING BROWNING OF APPLE SLICES DURING INFRARED DRYING Yashaswini S.S., Arun Kumar T. V., Pramod P. Aradwad, Indra Mani, Shalini G. Rudra and Bishal Gurung	27
26.	HYDROGEN STORAGE CHARACTERISTICS OF HIERARCHICAL CARBON DERIVED FROM BIOMASS USING THERMOCHEMICAL TREATMENT Bhupendra Ghritalahre, V.K. Bhargava and S. Gangil	28
27.	MANAGEMENT OF JACKFRUIT STEM BORER, BATOCERA RUFOMACULATA DE GEER (CERAMBYCIDAE: COLEOPTERA) Doddabasappa B., Mutthuraju, G. P. and Amarananjundeswara H.	29
28.	EFFECT OF IRON AND ZINC OXIDE NANOPARTICLES SYNTHESIZED VIA GREEN ROUTE ON GRAPES: AN ECOFRIENDLY APPROACH Yukti Verma and Ajay Kumar Upadhyay	30

Sl. No.	Title & Author of the Abstract	Page No.
29.	THERMAL DEGRADATION KINETICS OF PIGEON PEA STALK Parmanand Sahu, Sandip Gangil, Vinod Kumar Bhargav and Mattaparthi Lakshmi Durga	31
30.	ENVIRONMENTAL STRESS CONTRIBUTES TO AGE-RELATED HORMONAL DYSFUNCTION: POLYAMINES CAN CURTAIL NEURODEGENERATION, MAINTAINING FUNCTIONAL INTEGRITY OF NEUROENDOCRINE SYSTEM Nayan Anand Mate and Arnab Banerjee	32
31.	VERMICOMPOST INFLUENCING THE AGRICULTURAL YIELD & QUALITY Vrushali R. Mahala	33
32.	EFFECT OF DOPANTS (CE AND N) ON PHOTOCATALYTIC EFFICIENCY OF TiO₂ FOR DEGRADATION OF DICHLORVOS Vandana Yadav, Raj Kumar Singh and Vipin K. Saini	34
33.	A STUDY ON CHEMICAL AND STRUCTURAL CHANGES OF MICROPLASTICS INDUCED BY ADVANCED OXIDATION PROCESSES Aiswriya V P, Muthukumar Muthuchamy and Anbazhagi Muthukumar	35
34.	DEVELOPMENT OF AN ULTRASENSITIVE LUMINESCENT PAPER BIOSENSOR FOR QUANTIFICATION OF ARSENIC AND CADMIUM HEAVY METAL CONTAMINATION IN DRINKING WATER AND SEA FOOD Manisha Tiwari	36
35.	IMPACT ASSESSMENT OF KOTA STONE MINE SPOIL ON THE GROUNDWATER QUALITY IN THE REGIONS OF SOUTH-EASTERN, RAJASTHAN Gulshan Kumar Sharma, Gopal Lal Meena, Shakir Ali, Kuldeep Kumar and Ashok Kumar	37
36.	ESSENTIAL MINERAL AND HEAVY METALS AVAILABILITY IN VARIOUS WATER RESOURCES OF COLD DESERT HIGH-ALTITUDE REGION OF LADAKH, INDIA Vijay K. Bharti, Guru Charan, Arup Giri, Sudhir Kumar and OP Chaurasia	38
Technical Session 2		
37.	AGRO – PASTORALISM IN LADAKH: CURRENT TRENDS AND CHALLENGES Mohd. Raza, G.S. Rawat and Tsewang Namgail	41
38.	BAMBOO: CARBON SEQUESTRATION POTENTIAL Sunandini, Bhupender Dutt, Sakshi Singh, Reva Jaryal and Divesh Thakur	42
39.	BIOTECHNOLOGY'S POTENTIAL CONTRIBUTION TO SUSTAINABILITY OF WOOD PRODUCTS Akshay Pingale and Bhupender Dutt	43
40.	RENEWABLE SOURCED MULCH MATERIALS FOR SUSTAINABLE AND IMPROVED PRODUCTION OF STRAWBERRY CROP Nilimesh Mridha, Dipak Nayek, Ashok Yadav, Sourav Paul, Atul Singha, Manjunatha BS, Manik Bhowmick, H Baite, Biplob Saha, Surajit Sengupta, DP Ray, Sourav Pal, Amit Das, Leena Mishra and DB Shakyawar	44
41.	FISH SILAGE AS A FEEDSTOCK FOR THE GROWTH AND SIMULTANEOUS LIPID PRODUCTION BY THE OLEAGINOUS YEAST Shifa Naseema and Sunil Kumar Suman	45
42.	GREEN SYNTHESIS OF Ag/ZNO COATED ACTIVATED CARBON FOR THE REMOVAL OF Pb FROM AQUEOUS MEDIUM Asha Shende and Sandip Mondal	46
43.	IDENTIFICATION AND ANALYSIS OF MARINE LITTERS ON THE BEACHES OF GOA Kuldeep Singh, BK Ramprasad, Abhishek Chauhan, Jyoti Saini and Tanu Jindal	47

Sl. No.	Title & Author of the Abstract	Page No.
44.	ROLE OF LAND USE CHANGES IN URBAN WARMING AND IDENTIFICATION OF RESILIENCE MEASURES IN PERI-URBAN AREAS Jyoti Saini, Anamika Srivastava, Renu Dhupper, Kuldeep Singh, Anil Kumar Gupta and Bratati Dey	48
45.	RESPONSE OF SELECTED RANGE-RESTRICTED BIRD SPECIES OF WESTERN HIMALAYAN REGION TOWARDS CLIMATE CHANGE Garima Kumari, Parul Bhatt Kotiyal, Hukum Singh and Manoj Kumar	49
46.	TREND ANALYSIS OF METEOROLOGICAL VARIABLES OVER ALPINES IN CHAMBA DISTRICT OF HIMACHAL PRADESH, NW HIMALAYA, INDIA Vaneet Jishtu, Monika Chauhan, Brij Bhushan and G.S. Goraya	50
47.	BIOAUGMENTATION OF LOW-DENSITY POLYETHYLENE (LDPE)-CONTAMINATED SOIL WITH ENTEROBACTER CLOACAE AKS7: A NOVEL APPROACH TO PLASTIC WASTE MANAGEMENT Ranojit Kumar Sarker and Prosun Tribedi	51
48.	DETOXIFICATION OF PHENOLIC COMPOUND BY FUNGAL CRUDE ENZYMES Raja Singh Rawal and Sunil Kumar Suman	52
49.	ISOLATION OF COPPER-MINING BACTERIA FROM THE MUNICIPAL SOLID WASTE DUMPSITES OF KOCHI: A GREEN APPROACH FOR REMEDIATION Jyoti Varier, Lisa Jose and Sujatha C.H.	53
50.	MICROCRYSTALLINE CELLULOSE DECORATED WITH FE₃O₄ NANOCATALYST FOR THE MICROWAVE-ASSISTED SYNTHESIS OF THIOGLYOXAMIDES UNDER GREEN AND SUSTAINABLE CONDITIONS Vinay S Sharma, Achalkumar Ammathnadu Sudhakar, Pranav Shrivastav and Rajender S. Varma	54
51.	SELECTIVE REDUCTION OF ARYL CARBOXYLIC ACID USING AND SOLID ACID AND NABH₄ Siddharth S. Patel and Hitesh D. Patel	55
52.	XENOBIOTIC BIODEGRADATION POTENTIAL OF BREVI BACTERIUM SP. ISOLATED FROM COASTAL DUNES OF GOA: A GENOMIC APPROACH Dviti D. Mapari and Sandeep Garg	56
53.	AN OVERVIEW OF ASSESSING THE PERFLUOROALKYL COMPOUNDS IN WATER BODIES AND THE REMEDIATION METHODS FOR A SUSTAINABLE ENVIRONMENT Akhilghosh K A, Muthukumar Muthuchamy and Anbazhagi Muthukumar	57
54.	EXPLORING THE LARVICIDAL EFFECT OF CHEMICALLY SYNTHESIZED PRISTINE -MNO₂ NANOSTRUCTURES AND EVALUATING ITS TOXICITY USING OREOCHROMIS MOSSAMBICUS Shrikant R. Naik, Avelyno H. D'Costa, Siddhi. K. Salgaonkar, Antara Poi Raiturker Yuga Ghotge, Namita Naik, Roydnil Fernandes and Sheryanne Velho Pereira	58
55.	MAHANEEM BASED AGRO-FORESTRY SYSTEM: SUITABLE FOR CLIMATE SMART AGRICULTURE R.P.S. Deswal, N.K. Kaushik, M.K. Jat, Rameshwar Singh and Satyajeeet	59
56.	ROBUST ENVIRONMENTAL LAWS AND LEGAL INSTITUTIONS AND GREEN TECHNOLOGY FOR SUSTAINABLE DEVELOPMENT Dviti D. Mapari and Sandeep Garg	60
Technical Session 3		
57.	ARSENIC IN AGRICULTURE AND ITS MITIGATION TO PROTECT THE FOOD CHAIN Pankaj Kumar Srivastava	63
58.	CHARACTERIZATION AND CLASSIFICATION OF SOIL UNDER RAMIE GROWING AREAS OF FLOODED LOWE BRAHMAPUTRA VALLEY OF NORTH-EAST HIMALAYA FOR APPROPRIATE MANAGEMENT S. Ramachandran, R. K. Jena, P. Ray, Gulshan Kumar Sharma, K. K. Maurya, S. Hota, S. K. Singh and S. K. Ray	64

Sl. No.	Title & Author of the Abstract	Page No.
59.	FAMILY NET VESSEL COMPOST TECHNOLOGY: BEST WAY TO MANAGE KITCHEN WASTE Asha Sahu, MC Manna, Sudeshna Bhattacharjya, JK Thakur, A Mandal, AB Singh and AK Patra	65
60.	A STUDY ON THE VARIATION OF THE PHYSICOCHEMICAL, HYDRO BIOLOGICAL AND LIMNOLOGICAL PARAMETERS WITH RESPECTS TO THE EUGLENA BLOOM AND ANTHROPOGENIC FACTORS IN A HIGH, MODERATE AND LESS PRODUCTIVE POND OF BANKURA DISTRICT OF WEST BENGAL, INDIA Subhajit Mahanty, Keshab Chakraborty and Surajit Majumder	66
61.	ASSESSMENT AND IDENTIFICATION OF EXOPHYTIC AND ENDOPHYTIC BIOFLOC FORMING ORGANISM FROM SELECTED MARINE ALGAE SOURCED ALONG THE COASTAL AREA OF PANDURANGAPURAM, VISAKHAPATNAM Arnepalli Chandrasekhar, C. Manjulatha, Khadem Hussain Saeedi and D.V.V. Satynarayana Raju	67
62.	BENTHIC FORAMINIFERA, SEDIMENT CHARACTERISTICS AND ECOLOGICAL STUDIES IN THE NEAR SHORE ENVIRONMENTS OF PALK STRAIT, TAMILNADU, EAST COAST OF INDIA M. Suresh Gandhi, K. Kasilingam, Annmaria K. George and Gayathri G.S.	68
63.	CONSEQUENCES OF CLIMATE CHANGE ON THE ENDANGERED BATS OF WESTERN GHATS, INDIA Sreehari Raman, Shameer T.T. and Pooja Ushakumari	69
64.	CONSERVATION STRATEGIES OF RARE ETHNOMEDICINALLY IMPORTANT ISODON NILGHERRICUS BENTH H. HARA PLANT Godse N.H. and B.M. Gaykar	70
65.	DEATH BY NETS: MONITORING BIRD KILLS IN POMOGRANATE ORCHARDS Karthik N.J., Vijaya Kumara and Apoorva Kulkarni	71
66.	ECOSYSTEM SERVICES VALUATION OF A RAMSAR COASTAL LAKE IN KERALA WITH SPECIAL EMPHASIS ON WATER QUALITY AND LAND USE LAND COVER PATTERN Vishnu Sagar M.K, Sabu Joseph and Arunkumar P.S.	72
67.	INTERACTIONS BETWEEN INDIAN WILDLIFE AND HUMANS IN TIGER STATE (MADHYA PRADESH) Yogesh Singh and Pawan Shijoria	73
68.	NATURE-BASED SYSTEM FOR WASTEWATER TREATMENT AND REUSE Monali Muduli	74
69.	SPECTROPHOTOMETRIC MONITORING OF EXPLOSIVES WASTEWATER TREATED USING INTEGRATED TREATMENT METHODS Bhumika Sharma, S. Mary Celin, Garima Upreti and Ranu Gadi	75
70.	WASTEWATER REUSE IN AGRICULTURE Sumiran Dabruse and Abhipsa Khobragade	76
71.	DEVELOPMENT OF REVERSE TRANSCRIPTION LOOP MEDIATED ISOTHERMAL AMPLIFICATION (RT-LAMP) ASSAY AS A POINT-OF-CARE DIAGNOSTICS FOR SARS-COV-2 Shweta Kelkar and Rashmi H.M.	77
72.	LARGE SCALE SOIL MAPPING (1:10000) USING GEOSPATIAL TECHNIQUES – A CASE STUDY OF DAHOD DISTRICT, GUJARAT R. L. Meena, P.C. Moharana, Mahaveer Nogiya, R. S. Meena, S. K. Yadav, L. C. Malav, Brijesh Yadav, R. P. Sharma and B. L. Mina	78
73.	BIODIVERSITY STUDIES OF PIGMENTED BACTERIA AND FUNGI ISOLATED FROM PANJIM NALLAH WATER AND CHARACTERISATION OF SELECTED ISOLATES Marielou Ferrao	79

Sl. No.	Title & Author of the Abstract	Page No.
	Technical Session 4	
74.	ASSOCIATION OF ABO BLOOD GROUP PHENOTYPES WITH COVID-19: A CROSS-SECTIONAL STUDY IN GOA Neeha Sinai Borker and Nandini Vaz Fernandes	83
75.	ENDOPHYTIC FUNGI INOCULATION: A SUSTAINABLE WAY FOR INCREASING THE PHYTOCHEMICAL CONTENT IN STEVIA REBAUDIANA (BERTONI) BERTONI Rachna Verma and Reema Devi, Kiran	84
76.	SENSITIVITY OF ANDROGRAPHIS PANICULATA (BURM. F.) MORPHOTYPES TO SOIL MOISTURE STRESS: STATUS OF ANTIOXIDANT ENZYME ACTIVITIES AND ANDROGRAPHOLIDE YIELD Kuldeep Singh A. Kalariya, Deep Shahia, Parmeshwar Lal Sarana, Ram Prasanna Meena, Narendra Gajbhiyea, Rohan Sarkar and Geetha K.A.	85
77.	A NEW CALIX[4]PYRROLE FUNCTIONALIZED FLUORESCIN BASED SUPRAMOLECULAR COMPOUNDS FOR LIQUID CRYSTALLINE AND WINDOW LAYER SOLAR CELL APPLICATIONS Suryajit L. Rathod, Hitesh M. Parekh and Vinay S. Sharma	86
78.	EFFECT OF FENUGREEK SEED EXTRACT ON FLUORIDE INDUCED TESTICULAR DISORDERS IN ALBINO RATS Rajkumar Maiti and Debidas Ghosh	87
79.	SEMECARPUS ANACARDIUM LINN. LEAF EXTRACT EXHIBITS ANTI-BREAST CANCER PROPERTIES AND PROLONG THE LIFE-SPAN OF TUMOR-BEARING MICE Rajesh Kumar Singh, Bhagaban Mallik, Amit Ranjan and Anil Kumar Singh	88
80.	EVALUATION OF NEUROPHARMACOLOGICAL ACTIVITIES OF PERSIMMON (DIOSPYROS KAKI LINN.) FRUITS IN RATS Asmita Arondekar, Sweaton Fernandes, Nevita D. Rivonkar, Mithila R. Mhamal	89
81.	HYPERTENSION: A MAJOR THREAT TO HUMAN BEINGS Supriya Meena and Jaya Arora	90
82.	HYPO-TESTICULAR ACTIVITY ASSESSMENT OF HYDRO-METHANOL (60:40) SEED EXTRACT OF LUFFA ACUTANGULA IN HUMAN AND ALBINO RAT: AN IN-VITRO INVESTIGATION Tanusree Mondal and Debidas Ghosh	91
83.	INTERACTIVE EFFECT OF ELEVATED CO₂ AND VARYING NITROGEN REGIME ON THE PHYSIO-BIOCHEMICAL RESPONSE OF NEOLAMARCKIA CADAMBA Manish Singh and Hukum Singh	92
84.	NOVEL MECHANISTIC INSIGHTS ON MELATONIN MEDIATED PHYSIO BIOCHEMICAL, MICROSCOPIC, AND HISTOCHEMICAL MODIFICATIONS IN TOMATO PLANTS TO PROMOTE SALT STRESS TOLERANCE Tanveer Ahmad Khan and Qazi Fariduddin	93
85.	SEQUENCE ANALYSIS OF siet gene IN STAPHYLOCOCCUS PSEUDINTERMEDIUS ISOLATED FROM CANINE PYODERMA Prashantha M K; Shambulingappa B E; Sundareshan S; Kotresh A M; Rudresh B H; Madhavaprasad C B and Arun S J	94
86.	SIMULTANEOUS ESTIMATION OF METHOCARBAMOL AND IBUPROFEN IN FIXED DOSE COMBINATION BY UV SPECTROPHOTOMETRIC METHOD Premanand Velip, Anand Mahajan, Adison Fernandes and Shweta Borkar	95

Sl. No.	Title & Author of the Abstract	Page No.
87.	STUDIES ON CLINICAL SPECTRUM OF CANINE PYODERMA ASSOCIATED WITH STAPHYLOCOCCUS PSEUDINTERMEDIUS AND OTHER SECONDARY BACTERIAL INVADERS AND THEIR ANTIBIOGRAM PROFILE FOR EFFECTIVE THERAPY Prashantha M K; Shambulingappa B E; Sundareshan S; Kotresh A M; Rudresh B H; Madhavaprasad C B and Arun S J	96
88.	DEVELOPMENT OF LOW COST PISTON PRESS BRIQUETTING MACHINE M. Z. Sheikh, A. G. Mohod and Sakhi Katre	97
89.	DEVELOPMENT OF BIODEGRADABLE NANOCOMPOSITE PACKAGING MATERIAL FOR CHICKEN MEAT Kandeepan G., Monish. V., Sowndarya. A., Babji. Y., Spoorthi A., Srikanya and Ayesha A.	98
90.	DEVELOPMENT OF ELECTROCHEMICAL BASED IMMUNOSENSOR FOR THE DETECTION OF COLISTIN-A BANNED ANTIBIOTIC IN CHICKEN MEAT Harsh Kumar, Kamil Ku a and Dinesh Kumar	99
91.	DPP-IV INHIBITION (ANTI-DIABETIC) POTENTIAL OF KAPPA CASEIN HYDROLYSATES Harshita Sonarhi, Sathish Kumar M H and Rajani C S	100
92.	STABILITY ENHANCEMENT OF NATURAL FOOD COLOURS BY MODIFIED PROTEINS Charu Bisht, Neha Rawat and S.K. Sharma	101
93.	METABOLOME PROFILING OF COLOSTRUM, TRANSITION AND MATURE MILK IN MURRAH BUFFALOES Brijesh, Umesh Kumar, Monika Sodhi, Amarjeet, Divya Chanda, Abhishek, and Manishi Mukesh	102
94.	REACTIVE OXYGEN SPECIES AND DIABETES MELLITUS: ENVIRONMENTAL IMPACT Debidas Ghosh	103
95.	SIMULTANEOUS ENANTIOSEPARATION AND SIMULATION STUDIES OF ATENOLOL, METOPROLOL AND PROPRANOLOL ON CHIRALPAK IG COLUMN USING SUPERCRITICAL FLUID CHROMATOGRAPHY: A GREEN APPROACH IN SEPARATION SCIENCE Priyanka A. Shah, Pranav S. Shrivastav and Pranav Pandya	104
96.	STUDIES ON CLINICAL SPECTRUM OF CANINE PYODERMA ASSOCIATED WITH STAPHYLOCOCCUS PSEUDINTERMEDIUS AND OTHER SECONDARY BACTERIAL INVADERS AND THEIR ANTIBIOGRAM PROFILE FOR EFFECTIVE THERAPY Prashantha M K; Shambulingappa B E*; Sundareshan S; Kotresh A M; Rudresh B H; Madhavaprasad C B and Arun S J	105
97.	SIMULTANEOUS ESTIMATION OF METHOCARBAMOL AND IBUPROFEN IN FIXED DOSE COMBINATION BY UV SPECTROPHOTOMETRIC METHOD Premanand Velip, Anand Mahajan, Adison Fernandes and Shweta Borkar	106
98.	GREEN SYNTHESIS OF SILVER NANOPARTICLES MEDIATED BY GARCINIA INDICA (KOKUM) AND THEIR EFFECTIVENESS AGAINST ACNE-CAUSING BACTERIA Trelita de Sousa and Gautami S. Teli	107
99.	AMELIORATION OF ENVIRONMENTAL TOXICANT BENZO(A) PYRENE INDUCED TESTICULAR GAMETOGENIC AND STEROIDOGENIC DYSFUNCTIONS BY A NATURAL ARYL HYDROCARBON RECEPTOR ANTAGONIST RESVERATROL: INVOLVEMENT OF P38MAPK/ATF2/INOS SIGNALING Kuladip Jana, Bhaswati Banerjee, Satyajit Halder and Debidas Ghosh	108-109
100.	A NOVEL CALEBIN A ANALOG INDUCES ROS AND INHIBITS CANCER PROGRESSION VIA DISRUPTING PI3K/AKT PATHWAY Satyajit Halder, Pradip Shit, Ananya Dutta, Anup Kumar Misra and Kuladip Jana	110

Sl. No.	Title & Author of the Abstract	Page No.
101.	EVALUATION OF METAL TOLERANCE OF HALOPHILIC EUBACTERIAL AND ARCHAEAL STRAINS ISOLATED FROM SOLAR SALTERNS OF GOA INDIA Fazila Shaikh and Sanika Naik-Samant	111
102.	PERFORMANCE OF PLANT SPECIES ON THE TEXTILE EFFLUENT AFFECTED SOIL Dipak Kumar Gupta, Anil Kumar Shukla, Kamla K choudhary, SR Meena and Keerthika A.	112

Technical Session 1



International Conference on NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR SUSTAINABLE DEVELOPMENT (NTSD-2022)

30th November - 2nd December, 2022
Goa University, Goa

BIOMONITORING OF INDOOR AIR CONTAMINANTS USING SPIDER WEBS

Aradhana K S¹, M. Vishnu Sreejith, Usha K Aravind^{1*}

¹School of Environmental Studies
Cochin University of Science and Technology, Ernakulam, Kochi, Kerala, India

²School of Environmental Sciences
Mahatma Gandhi University (MGU), Kottayam, Kerala, India

ABSTRACT

The neurotoxic, immunotoxic, and other health problems associated with indoor air contaminants are of great concern. Hence, in recent decades there has been a substantial rise in studies pertaining to indoor air quality. The majority of the existing sampling techniques concerning indoor air monitoring involve the use of samplers made of synthetic materials that are less biodegradable. Thus, there is an urgent requirement to develop ecofriendly indoor air samplers that are cost effective and allows trace level detection of contaminants. The ability of spider webs to adsorb air pollutants makes it useful for measuring atmospheric pollution levels (Yu et al., 2007). In this context, the present work assessed the suitability of spider webs as a passive air sampler to determine the air contaminants in indoor spaces. The webs of Pholcidae species (a widespread spider throughout all indoor environments) were collected from three indoor work spaces. Heavy metal (As, Ba, Cd, Cr, Fe, Pb, Hg, Ni, Mn, and Zn) concentration in webs were monitored using inductively coupled plasma optical emission spectrometry (ICP-OES). Furthermore, the samples were subjected to ATR – FTIR spectroscopy. The results revealed the presence of various heavy metals, minerals and organic compounds in the web samples. The concentration of heavy metals in the samples followed the order Fe (14590 -24680 µg/g) > Zn (490-600 µg/g) > Hg (92-120 µg/g) > Cr (114.1-117.1 µg/g) > Cu (72-81.7 µg/g) > Pb (15.5- 36.9 µg/g) > Ba (7-8.7 µg/g) > As (BDL-1.2 µg/g) > Cd (0.4-1 µg/g). The IR spectra indicated the occurrence of quartz, calcite, and organic compounds having functional groups such as C-O, C-H and N-H (Figure.1) in the samples. This study demonstrated the use of spider webs as an excellent, eco-friendly and cost – effective passive sampler for monitoring indoor air contaminants at trace levels.




Figure 1: FTIR spectra of spider web

REFERENCE

Sreejith, M. V., Aradhana, K. S., Varsha, M., Cyrus, M. K., Aravindakumar, C. T., & Aravind, U. K. (2021). ATR-FTIR and LC-Q-ToF-MS analysis of indoor dust from different micro-environments located in a tropical metropolitan area. *Science of The Total Environment*, 783, 147066. <https://doi.org/10.1007/s00128-006-0917-y>

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**IMPACT OF ENVIRONMENT AND LIFE STYLE
ON GYNAECOLOGICAL CANCER RISK**

Nirmala K.* and Manjulatha C.

Department of Zoology, Andhra University, Visakhapatnam, India

ABSTRACT

The risk of gynaecological cancer (GC) is nonlinearly correlated with women's exposure to air pollutants (PM_{2.5}, CO, O₃, and SO₂). Residents in municipalities with greater levels of PM_{2.5}, a proxy measure of polycyclic aromatic hydrocarbons (PAH) and NO₂, had a higher risk of dying from ovarian cancer. The prevalence of pulp and paper manufacture was found to be highly linked with the incidence rates of ovarian cancer. Ovarian cancer risk factors include physical inactivity, obesity, talc-containing body powder used in the vaginal region, silica dust exposure, dry cleaning, telegraph and telephone work, and graphic and printing work. Women who had previously smoked cigarettes were more likely to develop cervical cancer as well as invasive and borderline mucinous ovarian tumors. Cervical dysplasia was more common in women in the highest residential benzene and diesel particulate matter (DPM) exposure categories and with high levels of home exposure to PAH. It was thought that exposure to PAHs influenced how cervical cancer developed and would strengthen the impact of high-risk human papillomavirus (HR-HPV) on cervical dysplasia. HPV infection and cervical, vaginal, and vulvar cancer have been linked in numerous studies. These comprised 62.5% of all cancer cases in the current analysis, and 76% of cervical cancer cases were found among low-income populations. Consumption of soybean milk, dietary additives, sweeteners, and preserved foods, exposure to plastics, cosmetics, and other pollutants like PM_{2.5}, could all be risk factors for uterine leiomyoma. Endometrial cancer risk factors include obesity, diabetes, and hypertension. Due to civilization, industry, and urbanization, humans are continually exposed to endocrine-disrupting substances (EDS), which promote the growth of endometrial cancer.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ASSESSMENT OF GROUNDWATER CONTAMINATION
IN COASTAL AQUIFERS IN NORTH GOA, GOA, INDIA**

Mahesh Mayekar¹, Trupti Asolkar² and A. Viegas¹

¹School of Earth, Ocean and Atmospheric Sciences, Goa University

²School of Biological Sciences and Biotechnology, Goa University

ABSTRACT

Fresh water is increasingly becoming a scarce commodity. Monitoring of fresh water resources is an important aspect considering the constant stress it faces due to ever increasing anthropogenic activities.

In this study, a network of 69 wells was selected from two contiguous coastal aquifers in Pernem and Bardez Talukas, North Goa, Goa and the data obtained was used to generate a groundwater flow-net. 18 wells were selected to determine their pH, electrical conductivity, total heterotrophic bacteria and coliform count. The pH ranges from 5.9 to 9.1, 15 samples are within the BIS (2012) drinking water standards, 2 samples being below and 1 above the permissible limit. Electrical Conductivity showed a range from 72 to 897 μ s/cm, 16 wells are within and 2 are above the BIS (2012) drinking water standards. The presence of indicator organism *Escherichia coli* was detected in all the 18 samples in the presumptive and completed tests of most probable number (MPN) raising serious concerns about the potability of the water.

The wells when superimposed on a groundwater flownet, indicates the contaminated wells and the areas where these contaminants are likely to migrate. Comparative analysis of coliform count depicted higher prevalence of coliform in Pernem Taluka as compared to Bardez Taluka. The high values of total heterotrophic bacteria and presumptive coliform count in certain areas may be ascribed to the close proximity of septic tank system with groundwater as well as its flow direction in these localities, which does not filter the contaminants infiltrating through the lithology, thereby creating conditions for contaminant to percolate into the groundwater.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**BASELINE PATTERN OF MICROPLASTICS, THE OMNIPRESENT
POLLUTANT IN THE WATERS OF VEMBANAD KOL WETLAND
OF KERALA, INDIA**

Nisari A R and Sujatha C H

Department of Chemical Oceanography, School of Marine Sciences
Cochin University of Science and Technology, Kochi, Kerala.

ABSTRACT

Multifunctional agrarian wetlands are contaminated with microplastics under the influence of intensive human activities and are considered to be a major sink and impacts the fragile systems. The degradation of larger plastic pieces to tiny pieces permits for extensive allocation and makes then persists in the aquatic food web. Attempts have been made to quantify the microplastic and the results showed that the fate and chemistry is directly linked and lethally affected the entire marine ecosystem. Hence the present investigation is initiated to evaluate the effect of these contaminants the identified hotspots. Here, the study is focused on the distribution microplastics in the waters of Kol wetland, a part of Vembanad Ramsar site and known the largest wetland in the south coast of India. To quantify the abundance of microplastics in the Kol wetland water, we samples were collected from monsoon season and subjected to microplastic extraction using density separation and filtration technique. The extracts were visually identified under a compound microscope and their chemical compositions were analysed using Fourier Transform Infrared (FTIR) spectroscopy. In all the examined water samples their presence were observed. Mean value of 30-70 numbers of microplastic particles in the experiment 2 liters of water were detected. Multiple types of microplastics representing fragments, films and fibers were detected. Polymers like polyethylene, polypropylene and polystyrene were conclusively identified with the analytical instrument. Filament shaped Polyethylene has been identified and is the most dominant types of polymer components in the analysed microplastics. The present investigation serve as the first baseline observation for evaluating the extent of microplastics in the water environmental matrices of Kol wetland and could extend for the effect of microplastics on aquatic biota.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**CONSTITUENTS OF MAJOR XENOBIOTICS IN THE
FISHERY HUB, VYPIN, KOCHI**

**Jayakrishnan Unnikrishnan², Lisa Jose¹, Emil John C.R¹,
Jyoti Varier¹, Nisari A.R¹, and Sujatha C.H^{1*}**

¹Department of Chemical Oceanography, School of Marine Sciences
Cochin University of Science and Technology, Kochi, Kerala

²St. Albert's College (Autonomous) Ernakulam, Kochi, Kerala

ABSTRACT

Estuarine contamination is a serious environmental threat which shows an increasing trend recently. Cochin estuary is one among them facing the same situation and Vypin zone is identified as one of the representative locales of the present investigation. The three surface sediment and surface water samples were taken from the Vypin area, which is an island region situated north side of the Cochin estuarine system and this fishing area is densely populated. Hence, the level of various xenobiotic constituents in the area needs to be studied and therefore analyzed. Influence of various anthropogenic activities leading to accumulation of both trace metal and various microplastic in the study site were assessed and evaluated. Analysis of trace metal in the sediment and water samples was done following the standard protocol and microplastic determination was done by density separation method. Grain size analysis was conducted by the pipette method. Total organic carbon, nutrients, chlorophyll, biochemical compositions of the sampled three surface sediment samples were analyzed in parallel following the conventional scientific method, along with microplastics and trace metals. The sediment texture in all three study samples was represented mostly by silt fraction, followed by sand and clay. A higher level of trace metals in this studied area along with various trans mended high amounts of microplastics were observed. These findings revealed to alter this vulnerable site and needs more research work to reduce the dangers of serious pollution in the whole aquatic ecosystem.

Keywords: Vypin fishing harbor, trace metals, Microplastics, Sediment and Water.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**CONTINUOUS MAPPING OF RIVERINE SYSTEM IN
KOCHI CITY USING MASS SPECTROMETRY**

Amitha Unnikrishnan¹, Usha K Aravind^{1*}, Shiny Thomas², and C. T. Aravindakumar³

^{1*}School of Environmental Studies, Cochin University of Science and Technology, Kochi, India

²SAIF-DST, Mahatma Gandhi University, Kottayam, Kerala, India

³School of Environmental Sciences, Mahatma Gandhi University, Kottayam, Kerala

ABSTRACT

The purity of river water is vital as it serves as a significant feeder to drinking water treatment plants for irrigation and other needs. However, one of today's most serious issues, particularly in developing nations, is river water pollution. It remains at the receiving ends of many effluent discharges, both point, and nonpoint sources. Though treated effluents are discharged, the screening studies indicate the presence of micropollutants. In the present study, the Periyar river, one of the significant rivers in Kerala that feed the entire Kochi city, is screened for the presence of micropollutants at selected sites. The study was carried out within a period of 2 years. The analysis of routine water quality parameters, ions, and organic contaminants was carried out from different sites continuously. The Total Organic Carbon (TOC) value at six sampling sites was observed to be higher than the allowed limit (< 4 mg/L). Site number 9 had the highest TOC reading (17.33 mg/L). The values of Chloride, Sodium, Potassium, Ammonia, and Magnesium ions were found to be above the allowed limits of the water quality standards. The values of basic water quality parameters were found to be within the allowed limit. The initial non-target analysis with LC-Q-ToF/MS showed the presence of 12 compounds, including mefenamic acid, metoprolol, lignocaine, and oxybenzone. To quantify their presence, target analysis was carried out. Prior to the analysis, SPE was used for the extraction, and eluents A and B were acetonitrile and water with 0.1% formic acid, respectively. Lignocaine and oxybenzone were below the quantification limit in 2018, but by 2020, their amount increased to 2.8 and 15.9 ng/L, respectively. Metoprolol concentration increased from 22.5 to 42.08 ng/L. While mefenamic acid concentration decreased from 680 to 0.78 ng/L. Currently, the number of study sites has increased, and the flow paths are identified to have more insights into the origin and occurrence of these micropollutants.

Keywords: Micro pollutants, Periyar River, (LC-Q-ToF/MS).

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**DISTRIBUTION PROFILE OF BIOCHEMICAL CONSTITUENTS IN THE
CORE SEDIMENTS OF VYPIN, THE FISHING HARBOUR OF KOCHI**

**Emil John C.R., Jayakrishnan Unnikrishnan, Lisa,
Nisari A.R., Jyoti Varrrior and Sujatha C H***

¹Cochin University of Science and Technology, Kochi, Kerala

²St. Albert's College (Autonomous), Kochi, Kerala

ABSTRACT

To understand the vertical fate and characterization of sedimentary organic matter composition in the major fishing harbour of Kochi, core sediments were collected from three selected sampling stations of Vypin region (V1, V2 and V3). This zone has been experiencing extreme adverse impact mainly from the industrial hub area, the Eloor, one of the world's 'top toxic hot spots' resides along the bank of Periyar. Sediment grain size was analysed by standard pipette method, while bulk organic matter content was determined by spectrophotometric methods. The sediment texture in both study areas was represented mostly by silt fraction, followed by clay and sand content. The present study revealed higher abundance of biochemical components in the Vypin region. High PRT/CHO ratios recorded at the surface layers of both V1 and V2 indicated fresh organic matter deposition in the study area and also lower chlorophyll-a/phaeophytin ratios of both V1 and V2 unveil anoxic conditions favourable for organic matter preservation. Thus, the results suggest that core sediments in the ecosystems tended to accumulate higher proportions of sedimentary organic matter by long period. Hence the present study revealed the significance of conservation ecosystems so as to increase fishing abundance, thereby reducing the pollution status and their role in influencing sedimentation and sustainability of the fishing zones of Kerala.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ESSENTIAL MINERAL AND HEAVY METALS AVAILABILITY
IN VARIOUS WATER RESOURCES OF COLD DESERT
HIGH-ALTITUDE REGION OF LADAKH, INDIA**

Vijay K. Bharti^{1*}, Guru Charan¹, Arup Giri¹, Sudhir Kumar² and OP Chaurasia¹

¹DRDO-Defence Institute of High Altitude Research (DIHAR), Leh, UT Ladakh, India

²Jaypee University of Information Technology, Wagnaghat, Solan, Himachal Pradesh, India

ABSTRACT

Water is an essential for all basic physiological functions of the body. The ground and surface water availability is limited in high-altitude region, including Ladakh, India. The availability of essential mineral nutrients and heavy metals in water resources has enormous impact on the health of growing plant, human, and animals. Therefore, evaluation of water availability and quality are extremely important for sustaining agriculture and animal husbandry practices along with maintaining human health. Therefore, the present study was conducted to evaluate concentrations of various essential minerals and heavy metals in ground (hand-pump), irrigation, pond and river water of Ladakh, India using ICP-OES spectrometry methodology. The results revealed that most of the essential minerals and heavy metals level in hand pump water were within the drinkable limit, whereas hand-pump, pond, irrigation, and river water from some site/place were high in Ca, Mn, Zn, As, Cd, Fe, Pb than the permissible limit of drinking water. Therefore, these water resources, if used for more extended periods without water treatment, may pose health-related issues to humans and animals from these elements. So, this study finding will help develop specific mitigation strategies for water management for drinking and other purposes in high-altitude region of Ladakh.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**PREVALENCE AND ASSESSMENT OF FLUORIDE LEVELS IN
GROUNDWATER OF HINGOLI, MAHARASHTRA**

Pushyamitra Joshi, N. G. Shinde and P. P. Joshi

Research Centre and Post Graduate Department of Zoology
K. J. Somaiya College of Arts, Commerce and Science, Kopergaon
Post Graduate Department of Zoology
Adarsh Education Society's Arts, Commerce and Science College, Hingoli (MS)

ABSTRACT

Water is vital component for life. As water is life, water parameters and quality of water is one of the challenges in today's scenario. Fluoride is one of the naturally occurring constituents of water. Fluoride is naturally occurring mineral which found in water. It often known as double edged weapon, as permissible limit is good for human's body but more than that are harmful for humans. Many health-related problems can be occurred due to high intake of fluoride dental fluorosis, skeletal fluorosis, osteoporosis, etc. Increasing concentration of fluoride levels in groundwater is one of major problem across the globe.

According to national drinking water mission's data, 20 states and 230 districts among them with 66 million peoples are endemic areas for fluoride related problems. Fluoride is often known as double aged weapon, as more than 1.5 mg/litre of fluoride is harmful for human body.

Groundwater is One of the important source of water in many parts of country. Especially in rural area, sometimes it is the only available source of water for primary use. Increasing concentration of fluoride in water due to anthropogenic and geogenic reasons is one of the international problems. Due to this, the present-day study conducted in District of Hingoli, Maharashtra. Analysis and assessment of groundwater samples for constituent of fluoride done as per the standard guidelines from world health organization (WHO) and American public health association (APHA). The results illustrate that some of water samples possess excess amount of fluoride. Present study reveals amount of fluoride from selected villages.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ADDRESSING EARLY BLOOMING IN ALMONDS UNDER
NORTH-WESTERN HIMALAYAN REGION OF KASHMIR**

Susheel Kumar Raina^{1,*}, Javed Iqbal Mir^{2,*}, Om Chand Sharma² and Sheikh M Sultan¹

¹ICAR-National Bureau of Plant Genetic Resources

Regional Station Srinagar, KD Farm, Old Air Field, Rangreth, Srinagar

²ICAR-Central Institute of Temperate Horticulture, KD Farm, Old Air Field, Rangreth, Srinagar

ABSTRACT

Almond is an important nut crop grown for its nutritious kernels. Almonds are rich source of protein, fibre, vitamin E, Omega-3 fatty acids and minerals like potassium, magnesium, phosphorus, calcium and iron. This nut crop blooms early in spring and blooming is often associated with early spring frosts. These frosts cause heavy damage to the flowers as well as discourage the activity of pollinators, leading to heavy reduction in fruit set, thereby posing serious economic loss to the growers. Late blooming almond cultivars escape early spring frosts and identification/development of late blooming almond cultivars has been an important breeding objective in this crop. To address the issue, we isolated *Flowering Locus C (FLC)* gene, a key gene associated with flowering, from almond cultivar Shalimar using degenerate primers. These primers amplified approximately 1 kb fragment revealing high sequence identity with *FLC* gene of *P. persica*. The coding region of the sequence (*PdFLC*) was submitted to Genbank under the accession number MG930898. This partial sequence of *FLC* gene from almond (*PdFLC*) was amplified later from several other cultivars and used for the construction of phenogram using Multalin software. The phenogram so created revealed three major groups viz. Group 1 including Primorskij, Makhdoom, Shalimar and California paper-shell; Group 2 including Pranyaj, Non-pareil and Merced; Group 3 including IXL, CITH4A, Waris and Drake. Moreover, transcriptome analysis of almond cultivars (Waris vs Ferralise) was done at flowering stage (pop corn stage of flowering) on an Illumina platform to identify the genes associated with blooming. Data processing employing pipeline: FASTQ, Trimmomatic, HISAT2, Stringtie, Stringtie merge and DESeq2 revealed prominent genes with significant up-regulation in local cultivar Waris as Ethylene Insensitive 3 while BADH acyltransferase and accelerated cell death were down-regulated.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

ANTI-DIABETIC PROPERTIES OF MARINE OLIGOSACCHARIDES

Devika Jadhav and Sanjeev C. Ghadi

School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau – Goa, India

ABSTRACT

The percentage of the diabetic population is increasing worldwide and is of great concern among children, and young adults. As per the International Diabetes Atlas, on an average every single Indian adult out of 11 is diagnosed by type 2 Diabetes mellitus. The risks and medical complications arising from diabetes are some of the major factors contributing to morbidity and mortality. Decreasing the mortality rates from non-communicable diseases is one of the sustainable development goals, adopted by the United Nations member states. Cases where patients have developed resistance to the available anti-diabetic drugs have been reported. Hence, there is a need to explore more molecules and compounds with potential anti-diabetic properties. In our study, we have isolated a Gram positive bacterial strain KK1, capable of enzymatically hydrolysing complex seaweed polysaccharides viz., agar, carrageenan, pectin and xylan. This is done by the machinery of a specialized set of proteins called carbohydrate active enzymes (CAZymes). The oligosaccharide hydrolysate produced by enzymatic degradation of respective polysaccharides was used for studying the anti-diabetic potential. These oligosaccharides were partially purified by precipitating the unhydrolyzed polysaccharides using ethanol and were further concentrated. The therapeutic potential of the oligosaccharides is known to vary concerning their source (parent polysaccharide), the enzymes used for hydrolysis, the degree of polymerization, and the presence of functional groups. These oligosaccharides demonstrated higher anti alpha-glucosidase activity in comparison to that of acarbose, a standard anti-diabetic drug. Thus, further exploring and bioprospecting of the anti-diabetic seaweed based oligosaccharides will help us get a step closer towards attaining the sustainable development goals.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**CHARACTERIZATION OF SEAWEED POLYSACCHARIDE DEGRADING
BACTERIA AND THEIR EXPLOITATION FOR ANTI-DIABETIC ACTIVITY**

Elaine Antoinette Da Costa

Department of Biotechnology, Goa University, Goa

ABSTRACT

Anthropogenic activities are the foremost cause of coastal eutrophication. A global rise in temperatures has caused algal and seaweed blooms that can disrupt the marine ecosystem. The bacterial degradation of seaweed biomass and the subsequent use of degraded oligosaccharide products for applications in agriculture, food and medicine needs to be explored. Polysaccharide-degrading bacteria from marine ecosystems are one of the major sources of Carbohydrate-Active enzymes (CAZymes). These CAZymes can be further exploited for the production of bioactive oligosaccharides. Several culturable polysaccharide-degrading bacteria were isolated from the coastal water samples where seaweed growth was prominent. They were isolated on a seawater-based medium with either agar, alginate, or carrageenan as the sole carbon source. The polysaccharide degradative activity was detected by the dye-based plate assay method. The polysaccharide hydrolysis was visualized as clear zones around the colonies, on flooding the plates with Lugol's Iodine for agarolytic and carrageenolytic activity. Alginate degradation activity was detected by flooding the alginate plates with absolute ethanol. Most bacterial strains degraded more than one polysaccharide, with most of them showing agarolytic and carrageenolytic activity compared to alginolytic activity. Bacterial strain AN104 was used to hydrolyse the seaweed polysaccharide carrageenan to produce a mixture of carrageenan oligosaccharides. These oligosaccharides were partially purified by ethanol precipitation followed by concentration by rotary evaporator. The concentration of oligosaccharides was estimated by the Nelson-Somogyi method. To determine the anti-diabetic potential of the carrageenan oligosaccharides, the anti- α -glucosidase activity was determined in comparison to the standard anti-diabetic drug acarbose. Results based on the IC50 values obtained reveal that the partially purified carrageenan oligosaccharides demonstrated a potentially much higher anti-diabetic activity as compared to acarbose.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ECO-FRIENDLY APPROACH FOR DETOXIFICATION OF A
HARMFUL TEXTILE DYE AND ITS PHYTOTOXIC EVALUATION**

Priyanka Bhagwan Jadhavar

Solapur University, Solapur, Maharashtra

ABSTRACT

The water is an essential resource for life on earth and for human development. The textile industry is one of the anthropogenic activity that most consume water and pollute water bodies also. The textile dyes significantly compromises the aesthetic quality of water bodies by increasing in BOD & COD, impair photosynthesis, inhibit plant growth, enter the food chain, provide recalcitrance and bioaccumulation, and may promote toxicity, mutagenicity, and carcinogenicity. Detoxification of Azo dye red 81 is selected for the present study because it is harmful if swallowed, May cause stomach discomfort, respiratory irritation, irritation to eyes, skin irritation in sensitive individuals, possible risks of irreversible effects. Particularly textile dye wastes are harmful to Agricultural and Aquatic habitat. Bioremediation is a pollution control technology that uses naturally capacity of microorganisms to break down toxic substances into non-toxic substances. The advantages of using microbes for bioremediation include; environmental friendly, their natural occurrence, ease of manipulation, high adaptive capacity, cheap production, fast growth rate, easy availability to treat large volumes of wastewater. In present study, in all total 05 dye decolorizing organisms were obtained. Out of these, DR-11 was efficient dye decolourizer and could decolorize the dye 98.00% in Nutrient broth having 10% NaCl. The potential strain was phylogenetically identified. The newly reported strain is most effective in decolourization, degradation and detoxification of Reactive Red 81 and having enormous potential of dye degradation under saline environmental conditions. The Microbial toxicity results proved that the degradation products of the dye were non-toxic to the agriculturally important microorganisms Viz. *Azotobacter spp.*, *Rhizobium spp.* and *Pseudomonas spp.* The phytotoxicity study reveals that the degraded products of the dye were non-toxic to the agriculturally important plant like *Triticum sp.* (Wheat). The use of such bacterium would be Very Cost-Effective and Eco-friendly technologies to achieve Sustainable Cleaner Production.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ELEVATED CO₂ AFFECTS DNA METHYLATION PATTERN
OF LOW AND HIGH ELEVATION POPULATIONS
OF *ARABIDOPSIS THALIANA***

**Ashwani Kumar Verma, Akanksha Singh, Rajneesh Singh, Prabhatam Mishra,
Shiv Narayan, Vivek Pandey, Pramod Arvind Shirke and Sribash Roy***

CSIR-National Botanical Research Institute, Lucknow, U.P.

ABSTRACT

Since the industrial revolution, atmospheric [CO₂] has risen from 280 ppm to over 410 ppm. It is expected to further rise to 550–700 ppm by 2050 and 650–1200 ppm by 2100, leading to global mean air temperature increase between 1.0°C to 3.7°C. Rise in atmospheric [CO₂] and temperature are imposing a great threat to the environment and in particular to plants because of their sessile nature. Understanding the molecular mechanisms of plant response to rising [CO₂] is critical for crop performance and the conservation of landscapes. However, plants responses towards rising atmospheric [CO₂] have largely been explored in genetical and morpho-physiological contexts. The epigenetic factors including DNA methylation which play critical roles in adaptation are largely unexplored. We have investigated the methylome, transcriptome and morpho-physiological responses of the two *Arabidopsis thaliana* populations evolved at high (3400 m amsl) and low elevation (700 m amsl) zones to elevated [CO₂] (e[CO₂]). We show that depending on the origin of the population, there is local level variations in DNA methylation when exposed to e[CO₂], but global methyl cytosine (mC) content remained unchanged. Further, there was loss of methylation and more protein coding genes were differentially methylated than other genomic contexts in both the populations but more so in the low elevation one. The differentially methylated genes of the two populations belong to distinct functional categories. More number of genes related to methylation machinery was down-regulated in the high elevation population than the low elevation one. Although there was no correlation between methylation and gene expression at global scale but a few genes exhibited methylation dependent expression level. Finally, we validated the gain or loss of methylation due to e[CO₂] treatment. Overall, our data suggested the two populations responded differently towards e[CO₂] with respect to methylome remodeling, phenotypic and molecular plasticity. However, methylome remodelling and molecular plasticity were more prominent in the low elevation population. Understanding the evolution of epigenetic response towards e[CO₂] may help in future crop improvement strategies.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ELUCIDATION OF MICROBIAL SIDEROPHORES IN
SALT TOLERANT KORGUT RICE VARIETY OF GOA**

Savia Bernadette Botelho

School of Biological Sciences and Biotechnology, Botany Programme
Block B, Goa University, Taleigao Plateau, Goa, India

ABSTRACT

Siderophores have been long studied for their iron sequestration ability under iron starvation conditions. Siderophores (gk. sidero-iron, phores-carrier) are high-affinity ferric ion-specific chelators with a low molecular weight of less than 10 kda, produced by various micro-organisms like bacteria, fungi, and also by some plants. This study aims to check the gene expression responsible for siderophore production in the salt-tolerant Korgut rice plant indigenous to the state of Goa. Korgut is known for its growth in high-salinity conditions and its nutritional benefit. Various rhizospheric, as well as endophytic bacteria, have been studied to produce siderophores helping the plants in the uptake of iron for their metabolic processes. Bacterial siderophores have a higher affinity for Fe than phytosiderophores and can remove Fe from Fe³⁺-phytosiderophore complexes. The other theory that has been suggested is that this occurs through ligand exchanges. The plant-produced siderophores (phyto-siderophores) interact with the bacterial siderophores-Fe complex, initiating the ligand exchange reaction. The plants absorb Fe via the iron-transferred Phyto-siderophores. The most common forms of siderophores are catecholates (produced by bacteria), hydroxamates (produced by both bacteria and fungi), and carboxylates (mainly produced by bacteria like *Rhizobium*, *Staphylococcus* and fungi like *Mucorales*). Some organisms produce mixed **siderophores** containing both catecholate and hydroxamate groups, such as heterobactin produced by *Rhodococcus erythropolis*. Many siderophores are small peptides synthesized by nonribosomal peptide synthetases, multimodular enzymes producing peptide products with a particular sequence without an RNA template. The expression of most proteins required for siderophore biosynthesis is regulated by a global iron-binding repressor protein called Fur (ferric uptake regulator).

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ENZYMATIC HYDROLYSIS OF CASTOR OIL BY
A NEWLY ISOLATED MICROBIAL STRAIN**

Shikha Singh¹, Sumit Sharma¹, Saurabh Jyoti Sarma* and Satinder Kaur Brar²

¹Department of Biotechnology, Bennett University, Greater Noida, India

²Department of Civil Engineering, Lassonde School of Engineering
York University, North York (Toronto), Canada, M3J 1P3

ABSTRACT

India is the leading producer of castor oil in the world with a market share of around 85-90%. In India approximately, 0.8 million hectares of land is devoted to the cultivation of the castor crop and around 0.7 % of the Indian farmers are engaged in castor cultivation. Castor oil consisting of 90% ricinoleic acid is a valuable industrial feedstock. Considering these facts, BASF, Arkema, Jayant Agro Organics Ltd., and Solidaridad came forward with the world's first sustainable castor bean program "PRAGATI". PRAGATI was launched in 2016 in Gujrat to ensure the uninterrupted supply of castor oil. Castor oil has diverse industrial applications due to its unique composition of fatty acids. The hydrolysis of the oil results in the production of ricinoleic acid, which can be converted into a number of valuable bio-based products. The hydrolysis of the oil could be achieved via a chemical route or biological route. The chemical route has disadvantages like the synthesis of by-products, energy squandering, and can also result in polymerization reactions. The biological route has advantages like product specificity and disposal of less harmful chemicals in the surrounding environment. The hydrolysis of castor oil via lipase from a newly isolated strain of *Aspergillus* has been presented in the study. The lipase activity obtained for 4% castor oil was around 6.68 $\mu\text{mol/ml/min}$. The concentration of ricinoleic acid and rate of hydrolysis were also determined.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**EVALUATION OF METAL TOLERANCE OF HALOPHILIC
EUBACTERIAL AND ARCHAEAL STRAINS ISOLATED FROM
SOLAR SALTERNS OF GOA INDIA**

Fazila Shaikh^{1*} and Sanika Naik-Samant¹

Biotechnology Discipline
School of Biological Sciences & Biotechnology, Goa University

ABSTRACT

Marine ecosystem is continuously exposed to metal pollution thus enabling estuaries and solar salterns to serve as an effective sink for these metals, leading to their high accumulation through evaporation. Estuarine environments thus serve as a good source of metal tolerant microbes which also contribute towards the biogeochemical cycling of inorganic elements. In this study, the halophilic strains from solar salterns of Agardvado and Maina, Goa-India were obtained on NTYE and ZMA agar plates incorporated with 25 % crude salt. The halophilic isolates were characterized based on their Gram character; tolerance to salt, antibiotic sensitivity, response to sodium taurocholate, evaluation of the pigment extracted and also examined for the presence of glycerol diether moieties. The potential eubacterial and haloarchaeal strains were screened for their metal tolerance/resistance ability using metal salts of Cu, Ti and Ag at 0.5, 1, 1.5, 2 and 4 mM concentrations. Growth was observed by monitoring the absorbance for each of the culture for an incubation period of 5-7 days. Culture tolerating/resisting metal salt exhibited alteration in pigment intensity as well as appearance or disappearance of pigment peaks. This indicated the role of cell wall components of potential cultures in withstanding the metal ions. Further, the potential cultures can also be used in synthesis of metal nanoparticles of nanobiotechnological significance along with the bioremediation applicability in metal contaminated saline soils.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**INTEGRATED FABRICATION OF A BIO-BASED COMPOSITE BY LACTIC
ACID CONDENSATION WITH RICE STRAW/LIGNIN FOR MAKING
SUSTAINABLE GOODS AND UTILITY ITEMS**

Sumit Sharma, Shikha Singh and Saurabh Jyoti Sarma*

Department of Biotechnology, School of Engineering and Applied Sciences,
Bennett University, Greater Noida, India

ABSTRACT

Rice stubble burning and its management issues are still faced every year. The advancement of polymer-based composite materials towards sustainability is exploring day by day. The chemically derived composites use precursors that are mostly derived from fossil fuel sources. Here, an approach was implemented for utilizing majorly renewable materials for making bio-based composite. A bio-composite material was fabricated by using rice straw/lignin as an intermediary compound during the stage of catalytic condensation of lactic acid. The reaction was catalyzed by Sn/Zn (2-5%) based catalyst at a temperature range of 80-160 °C. The rice straw was mixed to interlink with the condensing structure. Biodiesel waste glycerol (5-10%) was used as an additive for increasing its flexibility toward molding into any form. Other strengthening additives such as silica, starch, and calcium were added at a certain ratio during solidification and molded in various frames to form stable and sustainable structures. This study shows the utilization of rice straw or lignin in bio-composite formation. It has potential applications for making furnishing items, automotive parts, household items, or other goods/utilities that can be an alternative for sustainably utilizing renewable resources.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**METAL-ORGANIC FRAMEWORK BASED ADSORBENTS FOR THE
REMOVAL OF DICLOFENAC FROM SYNTHETIC WATER**

Amrutha Asokana¹ and C.T. Aravindakumar^{1,2*}

¹Inter University Instrumentation Centre,

²School of Environmental Sciences

Mahatma Gandhi University, Kottayam, Kerala, India

ABSTRACT

The presence of pharmaceutical and personal care products (PPCPs) in aquatic matrix highly demands newer innovations in water treatment protocol. Diclofenac (DCF), a very common non-steroidal anti-inflammatory drug with widespread occurrence is selected here as a model compound to evaluate the effectiveness of Metal Organic Frameworks (MOFs) as an adsorbent. MOFs are porous organic-inorganic materials having high specific surface area. The presence of functional groups and tunable pore size makes them an excellent material for water purification. In the present work, NH₂-MIL-101 (Al) is synthesised by a solvothermal method and characterised by SEM analysis and FTIR spectroscopy. DCF can be successfully adsorbed using NH₂-MIL-101 (Al). The effect of adsorptive conditions on DCF removal was also examined. The percentage removal of DCF was evaluated using high performance liquid chromatography (HPLC). NH₂-MOF showed above 90% adsorption capacity in all the studied pH. Adsorption at pH 4, 5 and 6 showed above 95% removal and the removal efficiency improved with increase in the weight of MOF from 42 to 98%. The adsorption efficiency in the presence of other coexisting compounds (Beta blockers) and ions (anions and cations) were also studied and they did not interfere the DCF adsorption except CO₃²⁻ and Al₃⁺. From the adsorption isotherm study it is found that Langmuir model positively fit to the adsorption than Freundlich model.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**MICROPLASTIC TOXICITY: AN ANALYTICAL INSIGHT INTO
DETECTION METHODS AND MYCO-DEGRADATION PATHWAYS**

Swati Solanki¹ and Rachana Singh^{1*}

¹Amity Institute of Biotechnology
Amity University Uttar Pradesh, Noida, India

ABSTRACT

Microplastics (MPs) have aroused a great deal of concern Due to their persistence in ecosystems and lack of degradability. Long-term exposure to microplastics can cause chronic toxicity, including impaired reproduction and malnutrition, threatening biota and humans. Aquatic populations that consume microplastics run the risk of choking, entanglement, and ingestion. Therefore, it is essential to develop highly efficient methods to remove MPs from the environment. In this regard, using fungi for microplastic degradation is beneficial owing to its diverse nature and effective enzymatic system. Extracellular and intracellular enzymes in fungi break down plastic polymers into monomers, producing water and carbon dioxide in aerobic environments and methane in anaerobic environments as bi-products. Further, fungi also secrete hydrophobins (surface proteins) which serve as a crucial aid in the bioremediation process by promoting substrate mobility and bioavailability. Therefore, the present review provides an insight into the mechanism and general pathway of fungal-mediated microplastic degradation. Additionally, analytical techniques for the monitoring of MPs degradation along with the roadblocks and future perspectives have been also discussed. To build an efficient and long-lasting technique for its management, more research is required to fully perceive the underlying process of microplastic biodegradation in the environment using fungus.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**PHYSIOLOGICAL RESPONSE OF ECTOMYCORRHIZAL
FUNGUS *PISOLITHUS* SP. ISOLATED FROM MINING SITE
TO TEMPERATURE STRESS**

Manasi Kamat and Lakshangy Charya

School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa, 403206.

ABSTRACT

Ectomycorrhizal (ECM) fungi are mutualistic symbionts of majority of woody plants. They play key role in nutrient cycling, production of plant growth promoting substances and antimicrobials against plant pathogens and combating abiotic stresses. *Pisolithus*, traditionally used in textile dyeing, is a well-known ECM basidiomycete in forestry trees such as *Eucalyptus*, *Acacia* and *Pinus*. With stresses such as temperature, salinity and so on deciding the success of reforestation, ECM fungi have become important bioinoculants for forestry plants to thrive in harsh conditions. The aim of this study was to isolate *Pisolithus* from fruiting bodies and study the effect of temperature on axenic culture.

Young puffballs lacked distinction, having only unconsolidated peridioles while developing basidiocarps showed mosaic arrangement of unconsolidated, consolidated, young and mature peridioles exhibiting colors ranging from white, yellow to olive. Free basidiospores were contained in upper portion of mature puffball. Fully ripe basidiocarp was found to readily crumble due to discontinuous peridium. Isolate PtC1 was successfully purified from mature, intact basidiocarp and produced fuzzy, golden-yellow colony on Glucose Mineral Salt Media (GMSM) with golden-brown, extracellular, phenolic pigment.

When grown on solid and in liquid media, PtC1 was found to grow from 15 to 37 with maximum colony diameter and biomass at 25. However, at this temperature, it produced least amount of pigment. In contrast, at 15 and 37 where biomass reduced by 81% and 70% respectively, higher amounts of pigment were produced. Colonies formed at these temperatures were compact with dark hyphae in comparison with fuzzy, light golden colony at 25. Further, the slow-growing colonies of PtC1 exposed to temperature of 15 grew at normal rate upon incubation at room temperature. These observations indicate that pigmentation may be a stress response mechanism in *Pisolithus*.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**UTILIZATION OF AGRO-RESIDUE IN HIGH-PERFORMANCE
SUPERCAPACITORS: A BRIEF CRITICAL REVIEW**

Ravi Kumar Sahu*, S. Gangil and V.K. Bhargav

ICAR-Central Institute of Agricultural Engineering
Bhopal, Madhya Pradesh, India

ABSTRACT

Energy storage plays important role in achieving global energy security and sustainability. In today's world, electricity is an essential part of our lives. A variety of renewable energy sources are used to produce electricity, including wind, hydropower, solar power, biomass, and geothermal. The storage of electrical energy is very important and needed option to fulfil the requirement of energy security and sustainability. Now days, the storage of electricity is major challenge. This problem can be dealt by the use of energy storage system. There are various types of energy storage devices for electrochemical energy like batteries, fuel cell, capacitors and supercapacitor, etc. According to review of different researches based on energy density and power density, supercapacitors have highest energy density among all existing electrochemical energy storage systems. Supercapacitor is an electrochemical unit comprising of anode, cathode, electrolyte and a separator. The electrode of supercapacitor is made of activated carbon which can be made from different crop residue and waste after appropriate processing. Carbon suitable for supercapacitor should have high specific surface area as 2000-3000 m²/g and pore size nearly 2–5 nm. The researchers have also found that micropores (<2 nm) perform better than mesopores (2–50 nm) over the surface of electrode material in supercapacitor. Reviews state that the analyses of electrode have shown the specific energy as 1–10 Wh kg⁻¹ and specific power as 500–10000 W kg⁻¹. This paper discusses about the use of agro biomass for making the base material for formation of effective electrode for use in supercapacitors.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**CARABID BEETLES (COLEOPTERA: CARABIDAE)
IN THE RICE AGRO-ECOSYSTEM OF KERALA**

Ashly Kurian*¹ and Sabu K. Thomas²

¹Department of Zoology, St. Joseph's College (Autonomous)
Devagiri, Kozhikode, Kerala, India

²Department of Zoology, University of Calicut, Tenhipalam, Kerala, India

ABSTRACT

Carabid beetles (Ground beetles) are abundant and normally live on the soil surface and inhabit almost all terrestrial habitats. They are active predators of insects and weed seeds hence they are considered as good biological control agents in agricultural ecosystems. Carabid beetles have bioindicator value, as they are sensitive to toxic substances (pesticides, herbicides and industrial fertilizers), agricultural practices (fragmentation and land management) and changes in the biotic factors of the environment. Studies in south-east Asian rice agro-ecosystems have established that Carabid beetles predate on many rice insect pests and maintain an equilibrium by regulating their number, acting as a keystone taxon in the rice agro-ecosystem. However, there is no comprehensive data about the Carabid beetles associated with the rice agro-ecosystem from Indian subcontinent and the major reason being the difficulty with the taxonomic verification of the family. The present study provides data about the species composition and abundance of Carabid community in a rice agro-ecosystem in Kerala. Forty-four Carabid species belonging to eleven subfamilies were recorded and verified. The data will be useful to identify the Carabidae species in the rice agro-ecosystems and to understand their importance as a dominant predator in rice agro-ecosystems.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

EDIBLE FILM PACKAGING-A NOVEL APPROACH IN FOOD PACKAGING

Abhipsa Khobragade^{1*} and Sumiran Dabrase²

¹Department of Agriculture, Rabindranath Tagore University, Bhopal (MP)

²Department of Agriculture, R. B. College of Agriculture, Wardha (MH)

ABSTRACT

India generated 3.3 million metric tons of plastic waste in 2018, according to a study made by the Central Pollution Control Board. Around 43% of manufactured plastics are used for packaging purpose. Huge amount of energy is required to degrade and recycle the food packaging. The main goal of food packaging is to provide a practical means of protecting and delivering food goods. Single-use food packaging takes a huge toll on our environment. The convenience of food packaging is outweighed by the waste and pollution that the packaging leaves behind. Most plastics do not biodegrade. Hence, there comes an urgent need to develop the edible packing and films. An edible film is defined as a thin layer, which can be consumed, coated on a food or placed as a barrier between the food and the surrounding environment. The main functions of this type of packaging are edibility and biodegradability. These natural films and coating are extracted from plants and animals in the form of polysaccharides, lipids, and proteins. The principle components for producing edible/biodegradable films are film forming biopolymers which includes the carbohydrates, proteins, solubilizing medium and plasticizers, etc.

On the other side, animals and humans are ingesting microplastic particles, via the food we eat and the water we drink. Thus edible film can both act as a barrier and protection, while enhancing quality and safety of food products. In the recent advances in edible film packaging bioactive peptides can play an important role, since they can fundamentally inhibit the growth of microorganisms by destroying their cell membrane. Edible films based on fruits and vegetables with novel health-promoting functionalities may also be developed, such as probiotic or prebiotic films, increasing their market appeal as healthy food components with desirable sensory properties.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**EFFECT OF NATURAL ANTI-BROWNING AGENTS IN AVOIDING
BROWNING OF APPLE SLICES DURING INFRARED DRYING**

**Yashaswini S.S.*, Arun Kumar T. V., Pramod P. Aradwad,
Indra Mani, Shalini G. Rudra and Bishal Gurung¹**

ICAR-Indian Agricultural Research Institute, New Delhi
¹ICAR-Indian Agricultural Statistics Research Institute, New Delhi

ABSTRACT

Apple is one of the most important fruit for a person's nutrition and vastly cultivated around the globe. The high moisture content makes them more susceptible to storage and transport. Hence, novel drying technology infrared drying is the energy efficient method and eco-friendly, adopted to avoid these losses. In apples polyphenol oxidase (PPO) causes enzymatic browning. It is well documented that the sample pre-treatment before drying can reduce various adverse phenotypic changes resulting from enzymatic reactions by inactivating the enzymes. Also, pre-treatment speeds up moisture evaporation and improves the color and texture of finished product to a large extent. Normally chemicals like KMS, ascorbic acid, cysteine, calcium salts used earlier but they affect the health of human beings. Hence, naturally available anti-browning agents like turmeric, honey and salt were used to prevent browning in apple slices before drying. The apple slices were soaked for 5, 10 and 15 min in naturally available anti-browning agents of three concentrations. The L, a and b values used to calculate browning index (BI) of pre-treated, infrared dried apple slices. The L value of turmeric water-soaked apple slices decreased as concentration increased and decreased with an increase in soaking time, a and b values increased as concentration and soaking time increased. In honey-treated apple slices, L value increased with concentration and decreased as soaking time increased, a and b values decreased as increased in concentrations and soaking time. Salt-treated apple slices shows increased L value and decreased a and b value with concentration and soaking time. The BI of control sample is 132.33 and turmeric, honey and salt treated slices ranged from 58.95 to 130, 50.60 to 34.39 and 28.53 to 46.53, respectively. The selected anti-browning agents were effective in controlling the enzymatic browning along with maintaining the quality properties of dehydrated apple chips.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**HYDROGEN STORAGE CHARACTERISTICS OF HIERARCHICAL CARBON
DERIVED FROM BIOMASS USING THERMOCHEMICAL TREATMENT**

Bhupendra Ghritalahre*, V.K. Bhargava and S. Gangil

ICAR-Central Institute of Agricultural Engineering
Bhopal, Madhya, Pradesh, India

ABSTRACT

An ever-increasing demand for hydrogen fuel has been a challenge for today's scientific workers. Production, transportation and hydrogen storage are the major challenges associated with H₂ fuel. Hydrogen fuel can be stored using different storage methods which have merits and demerits. Among all the storage methods i.e. pressurized gas, liquid nitrogen at cryogenic temperature, chemical method storage in metal hydride, storage of H₂ on carbon material is the safest and most reliable method. Many researchers utilize hierarchical carbon for electrical energy storage. Various literature reveal that carbon material for H₂ storage is an emerging technology. Hierarchical carbon having modified 3-dimensional structure and multimodal pore is prepared from different chemically or catalytically activation process. The output may reach up to the surface area of 1250-4000 m²/g and pore volume of 1.5 to 6 cm³/g. This level of surface area and pore volume show high potential for H₂ storage on carbon material. The percentage of hydrogen storage linearly increases with an increase in surface area. The H₂ storage on carbon having cellulose, hemicellulose and lignin ranges from 2.3 to 7.58 %wt. at 77K temperature and pressure of 10 Mpa. However, the variation in H₂ storage capability was observed when biomass was varied. Hydrogen storage on carbon is one of the prominent technologies for overcoming the challenges associated with the storage and transportation of H₂.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**MANAGEMENT OF JACKFRUIT STEM BORER, *BATOCERA
RUFOMACULATA* DE GEER (CERAMBYCIDAE: COLEOPTERA)**

Doddabasappa B.*, Mutthuraju, G. P. and Amarananjundeswara H.

Department of Entomology, Regional Horticultural Research and Extension
UHS Campus, GKVK, Bengaluru, KARNATAKA

ABSTRACT

The jackfruit is native to the rain forests of the Western Ghats of India and is cultivated throughout South and Southeast Asia, central and eastern Africa and Brazil. The study was conducted on effectiveness of different management practices of the trunk borer infestation in jackfruit garden, Tamaka, Kolar, Karnataka, India. The chemical and non-chemical treatments were imposed against jack stem borer infestation during 2017-2018 and 2018-2019. Sixty-five infested jack trees were selected and each treatment was applied for three infested trees. The observations on treatment effect were recorded at 15 days intervals from the date of treatments imposition. The efficacy of different treatments were grouped into four classes (I, II, III and IV) to represent the state of larval activity, with corresponding values (v) of 3, 2, 1 and 0 to calculate the degree of control (Sheng-ying *et al.*, 2009). Results from two years field experiment revealed that the jack stem borer can be effectively managed by placing the aluminium phosphide tablet (one tablet/ hole)+ sealed hole with Bordeaux paste ensured cent percent control of infestation. The next best management treatments are injection with DDVP @ 5ml/l+ sealed hole with Bordeaux paste (86.66%) and injection with Chlorantraniliprole @ 2ml/l+ sealed hole with Bordeaux paste (73.33%) followed by cent percent damage recovery and death of the tree noticed in untreated control.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**EFFECT OF IRON AND ZINC OXIDE NANOPARTICLES SYNTHESIZED VIA
GREEN ROUTE ON GRAPES: AN ECOFRIENDLY APPROACH**

Yukti Verma* and Ajay Kumar Upadhyay

ICAR-National Research Centre for Grapes, Pune, Maharashtra, India

ABSTRACT

For the past few decades, extensive research efforts have been made towards the preparation of cost-effective and eco-friendly nanostructured materials in the research fields of nanotechnology, a propitious branch of science that has received major success in the era of modern technology. Considering this, nanoparticles of iron and zinc were fabricated through novel green route. The obtained products (assumed as Fe-NPs and Zn-NPs) were subsequently characterized by Fourier Transform Infrared Spectroscopy, UV-vis spectra, Scanning Electron Microscopy and Particle Size Analysis techniques. The LCMS/MS was performed for the identification of biomolecules present in the grape pomace extract for formulating Fe-NPs and Zn-NPs. The morphology of Fe-NPs was monitored by SEM analysis and the particles were found in agglomerated form whereas the morphology of Zn-NPs was mostly spherical/agglomerated in shape.

To assess the impact of nanoparticles on grapes, field experiments were conducted. A significant increase in leaf Fe (+30.28% and +36.59%) and Zn content (+20.90% and +28.39%) was recorded after 14 and 28 days of application of Fe-NPs and Zn-NPs over conventional fertilizers in field studies. These findings suggested that Fe and Zn NPs have potential applications and could be used as a promising candidate for enhancing micronutrient content.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

THERMAL DEGRADATION KINETICS OF PIGEON PEA STALK

**Parmanand Sahu*, Sandip Gangil
Vinod Kumar Bhargav and Mattaparthi Lakshmi Durga**

Agricultural Energy and Power Division
ICAR-Central Institute of Agricultural Engineering, Bhopal, Madhya Pradesh, India

ABSTRACT

This study concentrates on pyrolytic kinetics and thermodynamics of intrinsic pseudo components of biomaterial. Thermogravimetric analyser was employed to investigate the thermal behaviour of pigeon pea stalk with four various heating rate (10, 20, 30 and 40 /min). Gaussian deconvolution function was used to resolve the individual spectrum of first derivative of conversion profile into three pseudo components (hemicellulose, cellulose and lignin). The Flynn-Wal-Ozawa (FWO), isoconversional method was used to estimate the pyrolytic kinetics of pseudo components. The activation energy for hemicellulose was computed high as 252-332 kJ/mol; whereas, for cellulose 128-145 kJ/mol and lignin 35-51 kJ/mol; for specific conversion level (0.1 - 0.9). The hemicellulosic conversion level $\alpha_H = 0.8$ showed highest activation energy as 332 kJ/mol. Also, the thermodynamic parameters; change in enthalpy (ΔH), Gibb's free energy (ΔG) and entropy (ΔS) for pseudo hemicellulose ($\Delta H = 278$ kJ/mol, $\Delta G = 178$ kJ/mol, $\Delta S = 169$ J/mol), pseudo cellulose ($\Delta H = 133$ kJ/mol, $\Delta G = 179$ kJ/mol, $\Delta S = -76$ J/mol) and pseudo lignin ($\Delta H = 36$ kJ/mol, $\Delta G = 232$ kJ/mol, $\Delta S = -260$ J/mol) were calculated. Presented article is an in-depth investigation to understand thermal degradation behaviour of pigeon pea stalk to obtain useful products.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ENVIRONMENTAL STRESS CONTRIBUTES TO AGE-RELATED
HORMONAL DISFUNCTION: POLYAMINES CAN CURTAIL
NEURODEGENERATION, MAINTAINING FUNCTIONAL
INTEGRITY OF NEUROENDOCRINE SYSTEM**

Nayan Anand Mate and Arnab Banerjee

Department of Biological Sciences, BITS Pilani, K K Birla Goa Campus

ABSTRACT

INTRODUCTION- Aging pathologies are greatly influenced by environmental stressors. Exposure to pesticides and environmental toxins are reported to contribute to neurodegeneration associated with both normal brain ageing and neurodegenerative disease. One of the inherent pathological features of aging evident very early on is the decline in reproductive function. Gonadotropin-releasing hormone (GnRH) is the master regulator of reproductive function. Polyamines are essential biomolecules that act in concert with neuroendocrine system. Previous studies have described protective functions of polyamines against diverse stresses. Polyamines act as reactive oxygen species (ROS) ROS scavenger, acid tolerance factor and chemical chaperone, and positive regulators for expression of stress response genes as well as modulator of neurohormones.

AIM- In the present study we investigated the protein and mRNA levels of polyamine and their corresponding biosynthetic enzymes such as ornithine decarboxylase (ODC), spermidine synthase (SRM) and spermine synthase (SMS) in the hypothalamus of female mice during ageing. Further we looked at the effect of polyamine on GnRH-I in presence/ absence of two important metabolic markers of ageing (which increase in response to environmental stress), TNF- α and lactate, through in vitro approach using GT1-7 neuronal cells.

KEY FINDINGS- Immunofluorescence study revealed that the expression of ODC1, SPM and SPD were relatively low to moderate in young and young adult hypothalamus; with a rise in intensity of expression in mature adult hypothalamus and then sharply declining in old hypothalamus. This pattern was almost similar to the expression of hypothalamic GnRH-I. TNF- α and lactate decreased hypothalamic GnRH I mRNA expression in GT 1-7 cells when treated for 24 hours. Polyamines, putrescine and agmatine, contrarily increased GnRH I mRNA expression significantly when treated alone for 24 hours to GT 1-7 cells. Also, polyamines increased GnRH I mRNA expression when treated in presence of TNF- α and lactate for 24 hours.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**VERMICOMPOST INFLUENCING THE
AGRICULTURAL YIELD & QUALITY**

Vrushali R. Mahala

Department of Forest Products
Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni-Solan (HP)

ABSTRACT

Vermicompost is a nutrient-rich, microbiological-active organic amendment that the interactions between earthworms and microorganisms during the breakdown of organic matter. It stabilized to divided peat-like material with a low C: N ratio, high porosity and high water holding capacity is the most nutrients are present in form to the plants. The agroforestry species produce huge amount of biomass and recycling of biomass by vermicomposting and other incorporation in soil by means of residue management practices for improvement of soil fertility. It prevents nutrient losses and increases the use efficiency of chemical fertilizers and it is free from pathogens, toxic elements, weed, seeds etc. it is necessary to monitor the temperatures of large-scale bin systems (high heat-retentive properties) as the raw materials or feedstocks used can compost and heating up the worm bins as they decay and killing the worms and the temperatures between 15-25°C (59-77°F). They can survive at 10°C (50°F) and above 30°C may harm them. The important of the agricultural sustainability for the future of the national economy and the Vermicompost is very valuable resource as organic fertilizers because it provides macro and micronutrients for plants and is a low cost, environmentally- friendly, invaluable alternative to chemical fertilizers. Earthworms have the potential to reduce the retention of pollutant and plant essential metals in the organic horizons by decomposing organic matter.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**EFFECT OF DOPANTS (CE AND N) ON PHOTOCATALYTIC
EFFICIENCY OF TiO₂ FOR DEGRADATION OF DICHLORVOS**

Vandana Yadav^{1*}, Raj Kumar Singh², Vipin K. Saini¹

¹Materials and Environmental Chemistry Research Laboratory
School of Environment & Natural Resources, Doon University, Dehradun, Uttarakhand, India

²Analytical Sciences Division, CSIR-Indian Institute of Petroleum, Dehradun, Uttarakhand

ABSTRACT

Dichlorvos (DHV) is a Class Ib (highly hazardous) insecticide, commonly used in developing countries, which is very harmful to aquatic organisms especially fish as it causes damage to the metabolism of the organism, which leads to their death. Therefore, this work is focused on the DHV degradation with Ce and N co-doped TiO₂ photocatalysts. Ce and N co-doped TiO₂ photocatalysts were prepared through sol-gel route by varying concentration of both Ce and N. The presence of CeO₂ cubic structure and rutile phase along with anatase phase was observed in the XRD pattern of all the co-doped samples. A significant reduction in surface area and pore volume has resulted due to Ce and N co-doping. UV-Vis DRS spectra exhibited a red-shift in the absorption edge due to synergistic effect of Ce and N dopants. The band gap values of co-doped samples are closer to nitrogen doped samples with lowest value of 2.27 eV. The XPS study proved the co-existence of Ce⁴⁺/Ce³⁺ via forming Ti-O-Ce type linkage. The formation of O-Ti-N bond linkage by N atom has been confirmed by FTIR and XPS results.

The photocatalytic performances of Ce and N doped and co-doped TiO₂ were investigated by conducting degradation experiments (under optimized conditions) of DHV in aqueous solutions. The results confirmed that photocatalytic activities of Ce-N-co-doped TiO₂ were higher for degradation of DHV as compared to pure and Ce or N singly doped TiO₂. This suggests that the presence of both the dopants (Ce and N) synergistically enhance the photocatalytic activity of co-doped TiO₂.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**A STUDY ON CHEMICAL AND STRUCTURAL CHANGES OF
MICROPLASTICS INDUCED BY ADVANCED OXIDATION PROCESSES**

Aiswriya V P¹, Muthukumar Muthuchamy¹, Anbazhagi Muthukumar^{1*}

¹Department of Environmental Science, School of Earth Science Systems
Central University of Kerala, Kasaragod, Kerala, India

ABSTRACT

Microplastics have gained global attention in recent years due to its increased prevalence in the body of living organisms including human beings and its possible ecological and toxicological impacts. Considering source reduction and clean up as the methods of MP pollution control, many physical, chemical and biological strategies have been progressively experimenting. Advanced oxidation process including photocatalysis, ozonation, electro-fenton and anodic oxidation is widely look over for its efficiency in degrading synthetic polymer particles leading to its mineralization. The present study investigated the efficiency of ozone in degrading virgin and processed polypropylene (PP) particles. Particles of size range 150 to 300 μm were dispersed in aquadest and subjected to ozonation under ambient temperature and pressure conditions with variable pH and reaction time. The experiment was conducted in a semi-batch reactor with varied pH conditions of 3, 5, 7, 9,11 and contact duration of 1, 2 and 3 hours. The extend of degradation was estimated by evaluating the gravimetric weight loss of the particles, COD analysis of solution, quantification of hydroxyl radicals and chemical structural changes of MP using Fourier Transform Infrared (FTIR) spectroscopy. Further analysis using SEM and XRD will be carried out to observe the morphological and structural changes of microplastics up on degradation.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**DEVELOPMENT OF AN ULTRASENSITIVE LUMINESCENT PAPER
BIOSENSOR FOR QUANTIFICATION OF ARSENIC AND CADMIUM
HEAVY METAL CONTAMINATION IN DRINKING WATER AND SEA FOOD**

Manisha Tiwari

School of Biological Sciences and Biotechnology, Goa University

ABSTRACT

Most of the Indian states are affected with heavy metal contamination mainly from mining, industrial effluents, improper industrial waste management and limited public awareness. In India, most of the rivers, estuaries and sea shores close to metropolitan cities are heavily contaminated. The effective way to reduce risks on common peoples is continuous monitoring of toxic metals in potable water or in the water body from where the sea food is regularly harvested or in the sea-food itself. Standard laboratory-based traditional analytical methods such as atomic absorption spectrometry (AAS) and atomic fluorescence spectrometry, inductively coupled plasma (ICP) are in practice for toxic metal quantification, but those traditional methods require expensive and bulky laboratory equipment, analytical expertise, sample transportation and pre-treatment.

Several research works have been reported for detection of heavy metal in drinking water and food utilizing chemically synthesized probe, engineered whole bacterial cell and genetically encoded probe. In most cases fluorescence probe was used which need light source, bleached easily, unstable, narrow detection limit and also require advanced instruments. Luminescence based sensor has capability to overcome with this problem. Here, I am developing low cost, ultra-sensitive, easy-to-implement and disposable heavy metal paper-based bioluminescence device using a brightest and smallest luminescent protein Nanoluc and the bacterial arsenic and cadmium metal regulatory trans protein and cis-factor. I am developing Nanoluc-trans factor recombinant proteins and immobilize oligomer cis-element on nitrocellulose paper. This sensor will be an intensimetric sensor, work with long pH range, independent to light excitation and capable to store at low temperature for long durability and cheaper.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**IMPACT ASSESSMENT OF KOTA STONE MINE SPOIL
ON THE GROUNDWATER QUALITY IN THE REGIONS
OF SOUTH-EASTERN, RAJASTHAN**

**Gulshan Kumar Sharma*, Gopal Lal Meena,
Shakir Ali, Kuldeep Kumar and Ashok Kumar**

ICAR-Indian Institute of Soil and Water Conservation, Research Centre, Kota, Rajasthan

ABSTRACT

Mining leads to the land degradation by stacking of mine waste, loss of top fertile soil and vegetation covers, abstraction and deterioration of natural drainage system, cutting of forest, groundwater depletion, and loss of plant and aquatic biodiversity and public health. A study has been conducted for assessing the impact of Kota stone mine on the groundwater quality. The groundwater quality parameters of 34 hand pumps used for drinking were analysed. The Fluoride (F) content in the groundwater were found to be ranged from 0.37-9.05 mg l⁻¹, and nearly 86.84 and 92.10% of the groundwater samples had F content higher than that of maximum desirable limit (WHO, 2017) and acceptable limit (BIS, 2012), respectively for drinking water (1.5 mg l⁻¹). The consumption of water having F⁻ content higher than 1.5 mg l⁻¹ usually causes dental fluorosis in both the children and adults. The TDS content in groundwater varies between 104 and 2240 mg l⁻¹, and 47.05% of the groundwater samples had TDS higher than 500 mg l⁻¹, which is not suitable for drinking purposes as per WHO and BIS. A similar trend as the TDS is recorded for the iron (Fe²⁺) content, 41.17% of samples were higher than 0.3 mg l⁻¹ as per WHO and BIS limit for drinking water (> 0.3 mg l⁻¹). Toxicity due to high concentration of the Fe in drinking water causes human health related hazards like hemochromatosis which results in organ damage, liver cirrhosis, hepatocellular carcinomas, fatigue, joint pain and hemosiderosis. The nitrate (NO₃⁻) content in 20.58% groundwater samples was found to be higher than 50 mg l⁻¹ as per WHO limit for drinking water. Consuming water having the NO₃⁻ content higher than 50 mg l⁻¹ may cause methemoglobinemia in the infants. In addition, the pH and the potassium content of Kota stone slurry were recorded 9.08 and 42 mg l⁻¹, respectively, which indicated the Kota stone slurry, is alkaline in nature and rich in the potassium. Results of analysis suggest that the Kota stone slurry could be utilized for reclamation of the acidic soils and also a suitable substitute for potassium fertilizer for the crop production in the region.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ESSENTIAL MINERAL AND HEAVY METALS AVAILABILITY IN
VARIOUS WATER RESOURCES OF COLD DESERT
HIGH-ALTITUDE REGION OF LADAKH, INDIA**

Vijay K. Bharti^{1,*}, Guru Charan¹, Arup Giri¹, Sudhir Kumar² and OP Chaurasia¹

¹DRDO-Defence Institute of High Altitude Research (DIHAR), Leh-194101, UT Ladakh, India

²Jaypee University of Information Technology, Wagnaghat, Solan, Himachal Pradesh, India

ABSTRACT

Water is an essential for all basic physiological functions of the body. The ground and surface water availability is limited in high-altitude region, including Ladakh, India. The availability of essential mineral nutrients and heavy metals in water resources has enormous impact on the health of growing plant, human, and animals. Therefore, evaluation of water availability and quality are extremely important for sustaining agriculture and animal husbandry practices along with maintaining human health. Therefore, the present study was conducted to evaluate concentrations of various essential minerals and heavy metals in ground (hand-pump), irrigation, pond and river water of Ladakh, India using ICP-OES spectrometry methodology. The results revealed that most of the essential minerals and heavy metals level in hand pump water were within the drinkable limit, whereas hand-pump, pond, irrigation, and river water from some site/place were high in Ca, Mn, Zn, As, Cd, Fe, Pb than the permissible limit of drinking water. Therefore, these water resources, if used for more extended periods without water treatment, may pose health-related issues to humans and animals from these elements. So, this study finding will help develop specific mitigation strategies for water management for drinking and other purposes in high-altitude region of Ladakh.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



Technical Session 2



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**AGRO – PASTORALISM IN LADAKH:
CURRENT TRENDS AND CHALLENGES**

Mohd. Raza^{1,2}, G.S. Rawat¹ and Tsewang Namgail²

¹Wildlife Institute of India, Dehradun, Uttarakhand
²Snow Leopard Conservancy India Trust, Leh, Ladakh

ABSTRACT

Agro – pastoral production system in Ladakh is mainstay of its economy and livelihood. However, due improved economic condition and infrastructure development such as roads, tourism and education it is undergoing changes. This study sought to understand current trend and challenges in sustaining agro – pastoralism in Kargil district. We used semi structured open – ended questionnaires per household to understand agriculture and livestock production system. Twenty village in five valleys comprising 519 households were covered. Overall, ten crops are cultivated in the region. Barley, alfalfa, wheat and peas are cultivated in all villages whereas oats is a new addition. Buckwheat, lentil, foxtail/proso millets and mustard cultivation is abandoned by most households in all the valleys. Likewise, five livestock types; sheep, goats, cow, yak, horse and donkey are reared. Overall livestock population is 5140 heads though it has decreased by 53 % (n=11054) in the aftermath of Kargil war in year 1999. In Chiktan Valley it has decreased by (71%; n=3216). In Drass and Suru Valley, it has decreased by 70%;2959 & 32%; 4879 respectively. Sheep and goats (shoats) from bulk of livestock population (n= 3766) whereas mule rearing is abandoned. Herd size of shoats per households has also decreased. Maximum decrease in herd size was observed in Chiktan Valley i.e., from 31.70 to 8.97 heads per households and least in Suru Valley i.e., from 14.88 to 10.23 heads. Similar declining trend is also observed for other livestock types. Overall lack of human resources due to off – farm income opportunities (53%) is a major challenge in sustaining agro – pastoral production systems, followed by fodder shortages (17%), climate (10%) and land division (8%). In order to sustain agro pastoral production system, it necessary to impart training in value addition of farm products in addition to opening market avenues for crops, handicraft and handloom products.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

BAMBOO: CARBON SEQUESTRATION POTENTIAL

Sunandini, Dr Bhupender Dutt, Sakshi Singh, Reva Jaryal and Divesh Thakur

Department of Forest Products, College of Forestry
Dr Yashwant Singh Parmar University of Horticulture and Forestry, Nauni-Solan (HP)

ABSTRACT

A continuous increase in the human population has generated huge demand for wood and wood-based products in service sector. To meet the industrial demand, large numbers of trees are cut down, causing serious climate change problems worldwide. The global temperature is rising with increase in carbon dioxide level in the atmosphere which was reached 414.7 ppm in 2021. To prevent irreversible disastrous effects on human society, the global temperature should be limited to 2 °C. At global level, the United Nations Framework Convention on Climate Change (UNFCCC) passed the resolution of the Paris agreement in 2015 to limit the rise of atmospheric temperature. Hence, for achieving sustainable development, non-woody forest timber, i.e., bamboo, has been gaining attention. In south-east Asia, over 1200 bamboo species are distributed in China, India and Myanmar. Asia accounts for approximately 65 percent of all bamboo-growing areas, with India reportedly home to 125 indigenous and 11 exotic bamboo species from 23 genera. Bamboo, being the fast growing species, has the potential to capture and store the carbon and restore degraded lands, which ultimately depends on its rate of growth and life cycle. The extensive root system of bamboo has been reported to improve soil properties in a short period of time. The carbon sequestration potential of bamboo varies from species to species. Due to its excellent qualities in physical and mechanical properties, bamboo is also considered as an alternative to wood.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**BIOTECHNOLOGY'S POTENTIAL CONTRIBUTION
TO SUSTAINABILITY OF WOOD PRODUCTS**

Akshay Pingale and Bhupender Dutt

Department of Forest Product
Dr Yashwant Singh Parmar University of Horticulture and forestry Nauni Solan HP

ABSTRACT

Wood is a naturally occurring, biodegradable, and renewable raw material that is used in building, as a feedstock in the paper and wood products industries, as well as in the manufacture of fuel. With the exception of the paper industry, biotechnology has traditionally received little attention in the wood product industries. However, biotechnology in forestry is also creating applications specific to forestry, such as genes for fiber modification, lignin reduction and extraction, and for the encouragement of straight stems and reduced branching. Legal constraints on conventional procedures have changed the situation as a result of growing environmental concern and scientific advancement. The goal of biotechnological wood protection strategies is to make wood more treatable with preservatives and replace chemicals with biological control agents. Utilizing fungi cultures and isolated fungal enzymes allows for the replacement of traditional chemical glues in the production of board materials. The rehabilitation of waste wood treated with preservatives benefits greatly from the use of biotechnology. Biotechnology will have many environmental and societal benefits. Faster-growing trees, developed through biotechnology, will contribute significantly to sustainable wood products by diminishing the demand for wood harvested from old growth and natural forest stands. Biotechnology can reduce the amount of lignin in trees intended for paper manufacture. Lignin which gives wood its strength must be removed in the pulping process. Trees that have less lignin or more-extractable lignin are more readily pulped, allowing mills to reduce the chemicals and energy required to purify cellulose (the basis for paper, packaging and many absorbent products) from wood. Thus, pulp mills are expected to better achieve their ambitious environmental objectives while reducing inputs and costs.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**RENEWABLE SOURCED MULCH MATERIALS FOR SUSTAINABLE AND
IMPROVED PRODUCTION OF STRAWBERRY CROP**

**Nilimesh Mridha^{1*}, Dipak Nayek², Ashok Yadav², Sourav Paul², Atul Singha¹,
Manjunatha BS¹, Manik Bhowmick¹, H Baite¹, Biplab Saha¹, Surajit Sengupta¹,
DP Ray¹, Sourav Pal¹, Amit Das¹, Leena Mishra¹ and DB Shakyawar¹**

¹ICAR-National Institute of Natural Fibre Engineering and Technology, Kolkata, India

²ICAR-Central Institute of Sub-tropical Horticulture (RS), Malda, India

ABSTRACT

Crop yields are generally limited by several factors like water, soil nutrients availability, proper managements etc. Strawberry is one of the most important soft fruits of the world after grapes and rich source of vitamin, iron and pectin content. It gives quickest return in shortest possible time. Besides antioxidant it contains phenols, flavonoids, dietary glutathionine which also exhibits a high level of antioxidant capacity against free radical species. Mulching manipulates the crop-growing environment to increase crop yield and improve product quality by controlling soil temperature, retaining soil moisture and reducing soil evaporation. A major limitation of polyethylene mulch involves disposal of mulch, if not properly disposed-off can fragment, and cause damage to environment i.e. land and water resources. The appropriate mulching materials, by controlling soil temperature and conserving soil moisture, can provide suitable soil microclimate for important crops (Chakraborty et al., 2008). Jute based renewable sourced non-woven mulch can be a sustainable alternative of synthetic mulch. The following study attempts to address the above issue.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**FISH SILAGE AS A FEEDSTOCK FOR THE GROWTH AND
SIMULTANEOUS LIPID PRODUCTION BY THE OLEAGINOUS YEAST**

Shifa Naseema and Sunil Kumar Suman*

¹Material Resource Efficiency Division
CSIR-Indian Institute of Petroleum, Haridwar Road, Dehradun, Uttarakhand, India.

²Academy of Scientific and Innovative Research (AcSIR)
CSIR-HRDC Campus, Ghaziabad, Uttar Pradesh, India

ABSTRACT

Extensive utilization of non-renewable energy sources and the pitfalls associated with the vegetable oils have led to several environmental concerns and political debates. Considering this, single cell oil produced by the oleaginous microorganisms has turned into a competitive alternative for biodiesel production. Under stress conditions and excess of carbon sources, oleaginous microbes can accumulate neutral lipids, mainly in the form of triglycerides (TAG). The present study proposes a novel feedstock derived from by-product of the fish processing industry (fish silage) for the growth and simultaneous lipid production by yeast strain *Rhodotorula mucilaginosa* IIP32. Under optimized culture conditions, maximum yeast biomass obtained was 6.5g/L. Lipid accumulated by the yeasts was in the range of 8-23% (w/w) and was initiated using in-situ transesterification process. Fatty acids from *R. mucilaginosa* IIP32 consisted majorly of Oleic acid 38%, Stearic acid 35% and Palmitic acid 14%, suggesting that the fatty acids could be used as a feedstock for biodiesel production. Utilization of fish silage besides providing economic benefits would also have a positive role in environment sustainability.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**GREEN SYNTHESIS OF AG/ZNO COATED ACTIVATED CARBON
FOR THE REMOVAL OF PB FROM AQUEOUS MEDIUM**

Asha Shende¹ and Sandip Mondal¹

Dept of Earth and Environmental Studies, NIT Durgapur, Durgapur, INDIA

ABSTRACT

In this work, green synthesis of Ag/ZnO nanocomposites on *Acacia Auriculiformis* (Akashmoni) Biochar to remove heavy metals from aqueous medium. SEM Analysis was done to verify the morphology and size of Ag/ZnO Nanocomposites whereas BET analysis for physical adsorption, XRD, and EDX techniques were used for crystallographic structure, chemical composition, and physical properties of the nanocomposites and FTIR for chemical properties respectively. The removal efficiency of the nanocomposites was determined by varying parameters like dosage of biochar (0.01g to 1g), initial concentration of heavy metal (1ppm to 100ppm), pH of the solution (2 to 8), and contact time between the adsorbent and the adsorbate (0.25min to 180min). Adsorption Kinetics study was used for investigating the mechanism of sorption using pseudo-first-order, pseudo-second-order, and Elovich kinetic models. The Adsorption Isotherm study was evaluated using Langmuir, Freundlich, Temkin, and Dubinin-Radushkevich isotherm models. This study also shows how Artificial Neural Network (ANN) was used for predicting performance of the adsorbent in the removal of heavy metals. Again, the parameters like dosage of biochar, initial concentration of heavy metal, pH of the solution, and contact time between the adsorbent and the adsorbate were used as input in the neural network. Working of the developed Artificial Neural Network model was determined using statistical measures like mean error, mean square error, root mean square error, and linear regression. The predicted ANN results and the experimental results were compared and a graph was plotted with experimented values on the X-Axis and ANN predicted values on the Y-axis. Ag/ZnO nanocomposite on biochar has been proved to be an effective and environmentally friendly adsorbent and can be utilized in various environmental applications.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**IDENTIFICATION AND ANALYSIS OF
MARINE LITTERS ON THE BEACHES OF GOA**

Kuldeep Singh*, BK Ramprasad, Abhishek Chauhan, Jyoti Saini and Tanu Jindal#

*#Amity University, Noida, Uttar Pradesh, India

ABSTRACT

Globally, the problem of marine litter raises concerns about environmental imbalance and its effects on coastal human life. The origins of marine litter include a variety of land-based and maritime activities. New initiatives in the marine sector, with a particular emphasis on coastal regions and thrust on the blue economy, show the government of India's vision for the nation very clearly. The UN Decade of Oceans Science for Sustainable Development (2021–2030) is in line with this.

A significant environmental concern has been the exponential rise in marine litter as a result of expanding marine activities. In order to evaluate the effectiveness of the system and establish policy measures to address the issue of marine litter, it is necessary to identify the sources of marine litter. Data on the beaches in Goa with random selection of the sample site was collected and analysis was carried to identify the major sources and major component of the marine litter on the beaches in Goa.

The study's mainly focuses on identification of the sources of marine litter to make effective policy decisions and implementation plans to address the problem of marine litter. The paper also aims to outline the main awareness-raising techniques against marine litter pollution.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ROLE OF LAND USE CHANGES IN URBAN WARMING AND
IDENTIFICATION OF RESILIENCE MEASURES IN PERI-URBAN AREAS**

**Jyoti Saini^{1*}, Anamika Srivastava¹, Renu Dhupper¹, Kuldeep Singh¹
Anil Kumar Gupta² and Bratati Dey³**

¹Amity University, Noida, Uttar Pradesh, India

²National Institute of Disaster Management, New Delhi, India

³Swastik Edustart, New Delhi, India

ABSTRACT

Changes in land use have emerged as one of the leading causes of climate change, or at the very least, they have intensified the process, as seen by the phenomenon of urban warming. Cities that were once thought to be environmentally comfortable are now referred to as heat islands during certain times of the year. Urban land is subject to changes because of pressure from human activities. A city's heat index will be higher if it has more "grey infrastructure," as concretization quickly absorbs and holds heat, warming the environment. Urban heat is mainly attributed to two factors. First, transmission lines bring tonnes of megawatts of electricity into cities, where it is converted into different forms of energy including heat and contribute towards formation of heat island (HI). Second, direct sunlight shining on grey infrastructure causes the HI effect.

An effort was made in this research to link the UHI status over urban grey infrastructure and green area for peri-urban areas (Dehradun and Gurgaon). LST maps for the years 2001 and 2021, respectively, were created using the thermal band of Landsat 5 and 8, respectively. Both ENVI and ArcGIS Desktop were used to complete the task. The generated data were then transferred to an excel spreadsheet and used for additional analysis. Maps, charts, and graphs were used to understand the results at the end. The quantity of grey infrastructure that is directly exposed to the sun, as well as the temperature gradient, have a significant impact on the city's heat index, according to the study's findings. Therefore, it is urgent to implement resilience mechanisms including the deployment of novel techniques to lessen direct solar exposure.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**RESPONSE OF SELECTED RANGE-RESTRICTED BIRD SPECIES OF
WESTERN HIMALAYAN REGION TOWARDS CLIMATE CHANGE**

Garima Kumari^{1*}, Parul Bhatt Kotiyal¹, Hukum Singh² and Manoj Kumar³

¹Forest Ecology and Climate Change Division, Forest Research Institute
Dehradun, Uttarakhand, India

²Genetics and Tree improvement Division, Forest Research Institute, Dehradun, Uttarakhand, India

³IT-GIS Discipline, Forest Research Institute, Dehradun 248006, Uttarakhand, India

ABSTRACT

Climate change, an alarming global phenomenon is considered a major threat to biodiversity. It is expected to alter the structure, function and composition of the Himalayan ecosystem which is believed to have an enormously beautiful aggregation of avifaunal species. These species are believed to face habitat loss due to an increase in temperature conditions, making a shift in their habitat. Therefore, predicting their distribution and suitable future habitats for them is the need of the hour to develop management options and conservation policies for adapting to changes in these escalating changing climatic conditions. An early initiative would help to understand the effect of climate change on species distribution. The study aims to develop an understanding of the distribution of selected bird species in the Western Himalayas in the current environment and to predict suitable future habitats for them under projected climate change scenarios. It will be observed whether future climate change and land use would modify the distribution of these species or not. Therefore, the species distribution modelling (SDM) approach using different RCPs will be used to account for different emissions and concentration trajectories. Likewise, the potential distribution of the selected bird species will be found. The models will be developed under the current climatic conditions and then projected onto various future climate change scenarios for different time frames in the future to monitor the changes in the distribution of the species and how the bird species will interact with their existing environment in the coming future. Then accordingly, with the help of results, conservation and management practices will be implemented.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**TREND ANALYSIS OF METEOROLOGICAL VARIABLES OVER ALPINES IN
CHAMBA DISTRICT OF HIMACHAL PRADESH, NW HIMALAYA, INDIA**

Vaneet Jishtu*¹, Monika Chauhan, Brij Bhushan and G.S. Goraya²

¹Forest Ecology and Climate Change Division,
Himalayan Forest Research Institute, Conifer Campus, Panthaghati, SHIMLA (HP) INDIA

²Himachal Pradesh Forest Department, Talland, Shimla (HP)

ABSTRACT

Different ecological studies conducted recently suggest that high altitude ecosystems are more vulnerable to fluctuations in meteorological variables than lower elevations. The floristic composition of high altitude is very rich and unique despite harsh environment conditions, which makes the flora sensitive to changes in external climatic variables. The increasing temperature trends reported in alpine regions have caused the species range shifts and increase in species richness, especially in deglaciated and permanent snow areas.

There is lack of sufficient ground-based data from these inaccessible areas. Thus, keeping this in view a study was conducted near Satrundi alpine area, located in district Chamba of Himachal Pradesh. Ambient temperature, relative humidity and dew point temperature data was recorded from the data logger installed at the study site using HOBOware. Data was analyzed and interesting results were obtained. The average ambient temperature recorded was 4.28°C, 4.51°C and 5.20°C with standard deviation of 6.67°C, 6.53°C and 5.74°C during 2019, 2020 and 2021 respectively. For relative humidity the average values recorded during study period were 67.29%, 64.49%, 62.76% with standard deviation 24.39%, 24.28% and 25.89% respectively. Data analysis of these meteorological parameters shows the significant variations over the years. The increasing trend of ambient temperature and decreasing trend of relative humidity (RH) is of great concern for climate induced environmental disasters.

As the unique and endangered biodiversity of alpine regions of Himalaya is under constant threat thus, anthropogenic activities responsible for degradation of these regions has also been discussed. This study also brought in focus the lack of sufficient ground-based data from these remote areas especially in Himachal Pradesh hence, more field studies are required to be carried out to know the actual effects and extend of climate change in order to suggest sustainable management strategies to save these sensitive regions.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**BIOAUGMENTATION OF LOW-DENSITY POLYETHYLENE (LDPE)-
CONTAMINATED SOIL WITH *ENTEROBACTER CLOACAE* AKS7:
A NOVEL APPROACH TO PLASTIC WASTE MANAGEMENT**

Ranojit Kumar Sarker and Prosun Tribedi

The Neotia University, Sarisha, West Bengal

ABSTRACT

An in-vitro study demonstrated that the soil bacterium *Enterobacter cloacae* AKS7, which breaks down low-density polyethylene (LDPE), accelerates the breakdown of UV-treated polymers. Its potential for contaminated LDPE habitat, however, has not yet been fully investigated. As a result, *in-situ* remediation was chosen for this investigation since it is more successful at getting rid of hazardous polymers. Four comparable soil-microcosm samples were similarly polluted with either UV-treated or untreated LDPE alone or in combination with AKS7 in order to study the same (T1 to T4). We detected considerable polymer degradation following the incubation in the bioaugmented soil contaminated with UV-untreated LDPE (T2). In addition, it was shown that UV-treated LDPE in AKS7 bioaugmented soil (T4) degraded polymers at a higher rate. Dehydrogenase and fluorescein di acetate (FDA) hydrolyzing activity were more active, which helped to explain why there were more microbes in the bioaugmented microcosm. The results of the carbon sources usage pattern of the BiOLOG Eco plate were used to assess the microbial functional diversity in order to comprehend the metabolic capability of the soil microcosms. Shannon-diversity index and Gini coefficient were shown to show the same, as functional-diversity include both microbial richness and evenness. According to the results, the microcosm that was exposed to UV-treated LDPE and challenged with AKS7 (T4) had the lowest Gini coefficient and highest Shannon-Weaver index when compared to the other microcosms. Overall, the findings showed that microcosm T4 had the greatest functional variety. When considered collectively, the findings showed that *Enterobacter cloacae* AKS7 might be employed as a bioaugmenting agent for the sustainable management of LDPE waste in contaminated sites without compromising the native soil microorganisms.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**DETOXIFICATION OF PHENOLIC COMPOUND
BY FUNGAL CRUDE ENZYMES**

Raja Singh Rawal^{1,2} and Sunil Kumar Suman^{1,2*}

¹Material Resource Efficiency Division

CSIR-Indian Institute of Petroleum, Haridwar Road, Dehradun, Uttarakhand, India

²Academy of Scientific and Innovative Research (AcSIR) CSIR-HRDC Campus, Ghaziabad, India

ABSTRACT

Phenolic compounds persistent in nature are a global concern, given their recalcitrant nature to degradation and adverse effects on humans and the environment. Removal of chlorinated phenols (CPs) from the environment usually involves expensive physical and chemical methods that produce harmful by-products; in contrast, the biological method (bioremediation) is an effective environment-friendly method without producing toxic by-products and is relatively cheaper. In the present work, the potential of ligninolytic enzymes such as laccase (Lacc) and manganese peroxidase (MnP) produced from *Pycnoporus* sp. (white-rot fungus) in the detoxification of 2-chlorinated phenol (2-CP) was studied. Both the enzymes exhibited stability in the pH range of 4-5 and temperature range of 50-65 °C. The lacc showed V_{max} and K_m values of 9.88 μMmin^{-1} and 41.91 μM with ABTS and 10.47 μMmin^{-1} and 93.56 μM with 2,6-DMP as substrate, whereas the MnP showed V_{max} and K_m value of 10.41 μMmin^{-1} and 7859 μM with guaiacol as substrate respectively. The crude enzyme cocktail detoxified the 2-CP (12.8 mgml^{-1}) into a less toxic form, confirmed by UV visible absorption spectra, HPLC analysis and plate toxicity assay. This study demonstrated that a crude enzyme cocktail produced by *Pycnoporus* sp. could potentially remove toxic phenolic compounds in more environmentally friendly ways.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ISOLATION OF COPPER-MINING BACTERIA FROM
THE MUNICIPAL SOLID WASTE DUMPSITES OF KOCHI:
A GREEN APPROACH FOR REMEDIATION**

Jyoti Varier, Lisa Jose and Sujatha C.H*

Department of Chemical Oceanography, School of Marine Sciences
Cochin University of Science and Technology, Kochi, Kerala

ABSTRACT

Municipal Solid Waste (MSW) dumpsites are the sinks of heterogenous refuses from different sectors of society. They also contain diverse indigenous microorganisms capable of withstanding these compounds, which can be exploited for various purposes. The leachate released from the MSW dumpsites presents potential risks to the surrounding environment. The dumpsite leachates constitute high concentrations of complex chemical pollutants such as ammonium, nitrates, pesticides, and trace metals. Among the trace metals, Cu (II) is one of the toxic metals serving as an essential micronutrient for plant growth. A reduced valence state of Cu (II) is Cu (0) having a wide application as heterogenous catalyst in the photoelectrochemical cells, sensors, inks, antimicrobial coatings and solar cells. As the conventional method for synthesizing the nanoparticles utilizes toxic chemicals and is costly, a sustainable method for reducing the toxicity of the Cu (II) and simultaneously extracting it for industrial application is the need of the hour. The exploitation of microorganisms for nanoparticle fabrication has garnered considerable global research interest. Among microbes, bacteria are an incredibly convenient target for green nanoparticle synthesis due to their variety and ability to adapt to different environmental conditions. It is known that individual bacteria are able to bind and concentrate dissolved metal ions and metalloids, thereby detoxifying their environment. Hereunder we report the green biosynthesis route for converting CuSO₄ ions into single-atom Cu (Cu₀) by an indigenous copper-resistant bacterium isolated from a major MSW dumpsite area of Kochi. The isolated bacteria were able to tolerate >1000 ppm of Cu. The bio transition of the Cu valence state was confirmed by various analytical methods (XPS, SEM-EDS and FT-IR). The results revealed that the bacterium was able to convert the Cu (II) (CuSO₄=100 mg/L) into Cu (0) within 24 hours. This microbial conversion is carried out naturally under aerobic conditions eliminating toxic solvents. These observations suggest and reveal that the MSW dumpsite bacterial population could naturally and efficiently extract metal ions such as copper and transform it into a highly valuable commodity in this new era of science.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



International Conference on
**NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

30th November - 2nd December, 2022
Goa University, Goa

**MICROCRYSTALLINE CELLULOSE DECORATED WITH Fe_3O_4
NANOCATALYST FOR THE MICROWAVE-ASSISTED SYNTHESIS OF
THIOGLYOXAMIDES UNDER GREEN AND SUSTAINABLE CONDITIONS**

Vinay S Sharma^{1*}, Achalkumar Ammathnadu Sudhakar²,
Pranav Shrivastav^{1*} and Rajender S. Varma^{3*}


¹Department of Chemistry, School of Sciences, Gujarat University, Ahmedabad, India.

²Departemnt of Chemistry, Indian Institute of Technology, Guwahati, Assam, India.

³Regional Centre of Advanced Technologies and Materials, Czech Advanced Technology and Research Institute, Palacký University in Olomouc, Šlechtitelů 27, 783 71 Olomouc, Czech Republic

ABSTRACT

Ferrite nanoparticles (Fe_3O_4 NPs) supported on microcrystalline cellulose (MCC), is introduced as an active heterogeneous catalyst for the preparation of the α -keto thioglyoxamides by α -heterofunctionalization of acetophenones with sulfur and secondary amines via a microwave (MW)-assisted protocol. This prepared catalyst shows consistent particle size of Fe_3O_4 NPs as characterized by PXRD, XPS, SEM, EDS, EDX, TEM, HR-TEM and FT-IR analysis. A wide variety of thioglyoxamindes are synthesized under optimized reaction condition to afford good yield of products in short reaction time. Importantly, the presented catalyst can be recycled for five successive runs with no major loss in its activity and is equally relevant for gram scale synthesis of thioglyoxamides. This environmentally benign approach offers several advantages and displays good green chemistry metrics (GCM), namely process mass intensity (PMI), E-factor and atom economy (AE). reaction mass efficiency (RME).



Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**SELECTIVE REDUCTION OF ARYL CARBOXYLIC
ACID USING AND SOLID ACID AND NaBH_4**

Siddharth S. Patel and Hitesh D. Patel*

Department of chemistry, Gujarat University
Navrangpura, Ahmedabad, India

ABSTRACT

Aryl alcohols and esters have immanent functionality in pharmaceuticals, agrochemicals, polymers, natural products, and food products. Also, these compounds can be employed as precursors for the preparation of various active pharmaceutical ingredients in pharma industries. The catalytic reduction of carboxylic acids into aryl alcohol is one of the most elementary and extensively employed transformations in synthetic organic chemistry. However, traditional methods (such as reduction using LiAlH_4) have their own drawbacks, such as the generation of inorganic waste, poor selectivity, lower yield. On the other hand, the catalytic hydrogenations of carboxylic acids were executed under high temperature and pressure largely to overcome their low reactivity and/or strong interactions with the metals catalysts. Thus, developing an alternative green protocol for chemoselective reduction of carboxylic acids to primary alcohols with wider functional group tolerance is still desirable. To this end, we have developed a new method for the reduction of aryl carboxylic acid by using recyclable solid acid CuPcS and NaBH_4 . These CuPcS and NaBH_4 system exhibit excellent yields and good selectivity, therefore providing alternative synthetic approaches for the conversion of carboxylic acids to primary alcohols with a wide range of functional group tolerance. The main advantage of both these methods includes many things, such as short reaction time, safer conditions, and an easy way of isolation and recycling of CuPcS.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**XENOBIOTIC BIODEGRADATION POTENTIAL OF BREVIBACTERIUM SP.
ISOLATED FROM COASTAL DUNES OF GOA: A GENOMIC APPROACH**

Dviti D. Mapari and Sandeep Garg*

Microbiology program, School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa, India

ABSTRACT

Xenobiotic compounds include drugs, pesticides, cosmetics, industrial chemicals, and environmental pollutants. Bioremediation strategies employing microorganisms is a promising solution to tackle environmental pollution caused by these compounds. Denovo whole genome sequencing of the halotolerant *Brevibacterium* sp. K11IcPPYGO002 and the pathway annotation of the predicted genes using KAAS (KEGG Automatic Annotation Server) revealed the presence of seventy-one genes responsible for the biodegradation of xenobiotic compounds. Analysis of the pathways revealed the presence of bacterial genes involved in the biodegradation of benzoate, aminobenzoate, fluorobenzoate, chloroalkane, chloroalkene, chlorocyclohexane and chlorobenzene, toluene, xylene, ethylbenzene, styrene, caprolactam, naphthalene, polycyclic aromatic hydrocarbon, steroid, and drugs. Genome mining of the *Brevibacterium* sp. K11IcPPYGO002 isolated from Coastal Sandunes of Goa is an effective practical approach for employing the isolate for the bioremediation of xenobiotic compounds.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**AN OVERVIEW OF ASSESSING THE PERFLUOROALKYL
COMPOUNDS IN WATER BODIES AND THE REMEDIATION
METHODS FOR A SUSTAINABLE ENVIRONMENT**

Akhilghosh K A¹, Muthukumar Muthuchamy¹, Anbazhagi Muthukumar^{1*}

¹Department of Environmental Science, School of Earth Science Systems
Central University of Kerala, Kasaragod, Kerala, India.

ABSTRACT

Chemicals in the per- and polyfluoroalkyl substances (PFAS) family exhibit strong fluorine and carbon linkages. They are more resistant to water and oil substances due to this quality. From their manufacturing through their consumption to their decomposition, these compounds are harmful to the environment. Furthermore, they endure extreme conditions and persist in the environment for a long time. More than 4500 PFAS and their precursors distributes in soil, water, and air after they released into the environment. It is in this specific context that environmental contamination due to PFAS should be addressed in the latest “Emerging Pollutants” category. Moreover, bioaccumulation of PFAS cause adverse health problems in plants, fishes and in mammals through food chain. Their uniqueness is that this new pollution can adversely affect the biotic and abiotic components, even in small quantities (one trillion parts). The scientific community has become profoundly aware of the adverse effects of these chemicals over the past couple of decades. Conventional wastewater treatment plants and methods followed by them are not an ideal solution for the removal of pfas from their sources. This study provides a critical analysis of the recent advancements in the detected amount of PFAS in waterbodies and the various economical and viable methods such as Advanced oxidation process (AOP) can be an effective technique for tackling the issues of PFAS.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



International Conference on
**NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

30th November - 2nd December, 2022
Goa University, Goa

**EXPLORING THE LARVICIDAL EFFECT OF CHEMICALLY SYNTHESIZED
PRISTINE α -MnO₂ NANOSTRUCTURES AND EVALUATING ITS TOXICITY
USING OREOCHROMIS MOSSAMBICUS**

**Shrikant R. Naik¹, Avelyno H. D'Costa², Siddhi. K. Salgaonkar,¹ Antara Poi Raiturker²,
Yuga Ghotge³, Namita Naik³, Roydnil Fernandes³ and Sheryanne Velho Pereira*³**

¹School of Chemical Sciences, Goa University

²School of Biological Sciences and Biotechnology, Goa University

³Department of Microbiology, St. Xavier's College Mapusa Bardez Goa

ABSTRACT

Mosquitoes are vectors of several diseases namely malaria, filariasis, chikungunya, yellow fever, lymphatic filariasis, Japanese encephalitis, Zika, West Nile and dengue. These diseases have been inflicting the lives of around millions of people annually. Indiscriminate usage of insecticides to curtail mosquito populations has resulted in devastating effects such as ecological imbalance, bioaccumulation, mammalian toxicity, and the dramatic emergence of resistant mosquito populations, to name a few. Hence there is a constant surge to find alternatives that are safe, target-specific, and equally effective or better. In the present study, we explore the larvicidal activity of α -MnO₂ nanostructures. X-ray Diffraction (XRD), Scanning Electron Microscopy (SEM), and Transmission Electron microscope (TEM) reveals that the MnO₂ nanocrystals have a hollandite (α -MnO₂) structure with a sea-urchin-like morphology and a high surface area. Larvicidal activity of MnO₂ nanoparticles was assessed against the larvae of *Culex quinquefasciatus* (*Cx. quinquefasciatus*), *Aedes aegypti* (*Ae. aegypti*), and *Anopheles stephensi* (*An. stephensi*), and revealed the toxicity in the order *Cx. quinquefasciatus* > *Ae. aegypti* > *An. stephensi* following the LC₅₀ values. Overall, this study highlights the potential of α -MnO₂ nanostructures as an effective mosquito control agent. Additionally, we also propose the use of the cichlid fish Tilapia (*Oreochromis mossambicus*) as an experimental model to study the toxic effects of these as-synthesised nanocrystals.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**MAHANEEM BASED AGRO-FORESTRY SYSTEM:
SUITABLE FOR CLIMATE SMART AGRICULTURE**

R.P.S. Deswal, N.K. Kaushik, M.K. Jat, Rameshwar Singh and Satyajeeet

CCS HAU, Regional Research Station, Bawal-123501, Haryana

ABSTRACT

A field study was carried out at research farm of CCS HAU, Regional Research Station, Bawal (Haryana) in rabi 2021-22 to study the effect of different densities of Mahaneem (*Ailanthus excelsa*) plantation on performance of wheat varieties grown as intercrop. The soil of experimental field was loamy sand in texture and alkaline in reaction (pH 8.22 to 8.30). The experiment was laid out in split plot design with three replications. *A. excelsa* was planted in September 2013 with four different tree densities viz. 200 (10 x 5m), 150 (10 x 6.5m), 100 (10 x 10m) and 50 (10 x 20m) trees/ha (spacing) and four varieties of Wheat (*Triticum aestivum*) i.e. WH-711, WH-1105, HD-2967 and HD-3086 were grown as intercrop. Recommended dose of fertilizers for the crop was applied as per packages practices of CCSHAU, Hisar. Irrigation and plant protection measures were applied as per recommended practices. Unusual high temperature was recorded in March, 2022. Results reveal that with decrease in tree spacing the yield of the wheat varieties increased significantly. All the wheat varieties recorded maximum yield under 10x5 m spacing. It has been recorded first time in Mahaneem based agroforestry system. It may be due to moderating effect of trees on high temperature in the month of March, 2022. The growth of trees in terms of height and girth at breast height (GBH) in sole plantation of Mahaneem as well as with intercropping did not differ significantly. However with close spacing / higher density of plantation tree height was slightly less and slight decrease in tree GBH was also recorded but statistically the difference was not significant. Maximum carbon sequestration was recorded under 10 x 5 m plantation of Mahaneem with wheat varieties followed by 10 x 6.5 m spacing. The soil organic carbon ranged from 0.19 to 0.22 percent, pH was recorded 8.19 to 8.21 and available phosphorus was recorded maximum in 10 x 5 and 10 x 6.5 m plantation after harvesting of wheat crop. Similarly, available K ranged from 166.35 kg/ha in 10 x 5 m plantation to 165.3 kg/ha in 10 x 20 m plantation.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ROBUST ENVIRONMENTAL LAWS AND LEGAL INSTITUTIONS AND
GREEN TECHNOLOGY FOR SUSTAINABLE DEVELOPMENT**

Dviti D. Mapari and Sandeep Garg*

Ravinder Kumar, Professor and Dean, Faculty of Social Sciences
Jamia Millia Islamia – NAAC A++ Accredited and NIRF 3rd Rank Central University, New Delhi

ABSTRACT

The fight for a clean environment continues in a climate of ever-increasing urgency. Constitution of India imposes a fundamental duty on every citizen (under Article 51A) to protect and improve the natural environment, including forests, lakes, 'rivers and wildlife', and to have compassion for living creatures. Although before India's independence in 1947, several environment legislations existed; the real impetus for bringing about a well-developed framework came only after the UN Conference on the Human Environment in Stockholm in 1972. Under the influence of this declaration, the National Council for Environmental Policy and Planning was set up within Department of Science and Technology. This Council later evolved into a full-fledged Ministry of Environment and Forests (MoEF) in 1985, the apex administrative body in the country for ensuring environmental protection. The Environment Protection Act (EPA), 1986, came into force soon after the infamous Bhopal Gas Tragedy. It is considered an umbrella legislation as it fills many gaps in the existing laws. It empowers Central Government to establish authorities equipped with the directive of preventing environmental pollution in all its forms and to tackle specific environmental problems, peculiar to different parts of the nation.

Sustainable development meets the needs of the present without compromising the ability of future generations to meet their own needs and provides a framework for humans to live and prosper in harmony with mother nature rather than, as we have for centuries, at nature's expense. However, sustainability does not seem to have an adequate legal foundation besides green technology, despite many environmental and natural resources laws (discussed above) in force. Hence, if we are to make significant progress toward a sustainable society, we will need to develop robust laws and legal institutions, green technology and ensure their effective and firm reinforcement by creating a strong will among the concerned.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



Technical Session
3



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ARSENIC IN AGRICULTURE AND ITS MITIGATION
TO PROTECT THE FOOD CHAIN**

Pankaj Kumar Srivastava

CSIR-NBRI, Lucknow, Uttar Pradesh

ABSTRACT

Arsenic (As) is a naturally occurring toxic metalloid at a mean content of 5 mg kg⁻¹ in the earth's crust. The potential risks of arsenic to ecosystems and human health, particularly in the South Asian region (India and Bangladesh, etc.), have been thoroughly investigated. Human health toxicity resulting from the use of arsenic-contaminated groundwater for drinking and irrigation is a global concern, particularly in Asia and Southeast Asia. Groundwater from tube wells used to irrigate agricultural fields is adding large amounts of arsenic to the soil every year, resulting in increased levels of arsenic in crops grown on arsenic-contaminated arable soils. Arsenic accumulates in the edible parts of plants and subsequently contaminates the food chain. It is highly warranted to initiate monitoring and assessment programmes for arsenic contamination in irrigation groundwater sources, agricultural fields being irrigated using those sources and the crops grown in those fields. Because data from spiked soil and hydroponics experiments cannot be extrapolated as truly representative of the field condition of arsenic contamination of soil and crops. An understanding of soil arsenic content and other soil properties in relation to its uptake and toxicity to crops is therefore urgently needed under field conditions. It is revealed that soil total arsenic content is not likely a good predictor of arsenic uptake and toxicity under different soil types and micro-climatic conditions. It is the bioavailable fraction of total arsenic in specific soil conditions, which is potentially causing a threat to the crops/plants. Identification of geographic areas either presently contaminated with arsenic or susceptible to its contamination, is an important step for risk assessment and developing remediation strategies. The mapping of arsenic content which is being transported from soil or irrigation water to edible plant parts is important for protecting against human arsenic exposure through arsenic contamination of the food chain. Assessment of soil arsenic bioavailability may greatly support the scope of remediation required at contaminated sites. It is concluded that future R&D should be focused on speciation, transformation and bioavailability of arsenic in soil and associated soil properties to reveal tangible latent hazards of soil arsenic contamination to agriculture and related food chain links.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**CHARACTERIZATION AND CLASSIFICATION OF SOIL UNDER RAMIE
GROWING AREAS OF FLOODED LOWER BRAHMAPUTRA VALLEY OF
NORTH-EAST HIMALAYA FOR APPROPRIATE MANAGEMENT**

**S. Ramachandran¹, R. K. Jena², P. Ray³, Gulshan Kumar Sharma⁴,
K. K. Maurya⁵, S. Hota⁵, S. K. Singh⁶ and S. K. Ray⁷**

¹ICAR-Indian Institute of Horticultural Research, Hessaraghatta Lake Post, Bengaluru, Karnataka

²ICAR-Indian Institute of Water Management, Bhubaneswar, Orissa

³ICAR-Indian Agricultural Research Institute, New Delhi-100012

⁴ICAR-Indian Institute of Soil & Water Conservation Institute, Dadwara, Kota, Rajasthan 324002

⁵ICAR- National Bureau of Soil Survey & Land Use Planning Regional Centre, Jorhat-
785004, Assam

⁶ICAR-Central Coastal Agricultural Research Institute, Goa – 403402.

⁷ICAR-National Bureau of Soil Survey & Land Use Planning, Kolkata- 700097

ABSTRACT

A study was conducted in the Ramie [*Boehmeria nivea* (L.) Gaud.], a growing area of Brahmaputra River derived soils in Barpeta district of Assam. The soils were studied to characterize and classify for appropriate crop management. The quality of crops is dependent on soil properties. Hence, the soil was studied and classified for crop suitability. For this purpose, eight representative profiles from different landform units of ramie growing area were characterized and classified. The Ramie growing soils are very deep, very dark grayish brown (10YR 4/2) to dark grayish brown in colour. The structure of the soil is granular to angular blocky in structure, loam to silty clay loam. The soils are strongly acidic to moderately alkaline in nature (pH 4.0 to 8.2), and the electrical conductivity varies from 0.001 to 0.003 dSm⁻¹. The OC is low to high in organic carbon (0.4 to 1.2 %). Anion (Al³⁺ and H⁺) activity ranges from 0 -1.63 and 0.2-1.59 respectively. Low cation exchange capacity (1.8 to 6.6 cmol (P⁺) kg⁻¹), calcium was the dominant cation, followed by magnesium, sodium and potassium. The base saturation percent was 27.64 to 96.36 percent. The fertility status of ramie growing soils was medium to high. Taxonomically, these soils were classified as *Typic Dystrudepts*, *Typic Fluvaquepts* and *Typic Udipsamments*. The soil-site suitability indicated that the *Typic Udipsamments* soils were marginally suitable, as *Typic Fluvaquepts* were moderately suitable and *Typic Dystrudepts* soils were highly suitable for ramie crop.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**FAMILY NET VESSEL COMPOST TECHNOLOGY:
BEST WAY TO MANAGE KITCHEN WASTE**

**Asha Sahu*, MC Manna, Sudeshna Bhattacharjya
JK Thakur, A Mandal, AB Singh and AK Patra**

ICAR- Indian Institute of Soil Science, Bhopal, India

ABSTRACT

In India rapid urbanization, escalating population and food consumption rate led to increase in city waste (CW) generation in India. Per capita generation of CW ranges from approximately 0.17 kg to 0.62 kg per person per day in small towns and cities respectively with a mean of 0.4 kg per capita. About 40% of CW is considered to be organic and biodegradable, of which only 14% (9.1 Mt) being composted. Organic CW (OCW) mainly comprised of discharges from household and can be better designated as Kitchen Waste (KW). The KW contains high moisture content and is characterised by about 60% carbohydrates, 20% protein and 10% lipid. Thus, a strong technology is needed to segregate KW at source level and convert it into stable and nutrient enrich compost. ICAR-Indian Institute of Soil Science, Bhopal, has developed Family Net Vessel Compost (FNVC) technology using three epigeic earthworms (*Eisenia fetida*, *Eudrilus eugeniae* and *Perionyx excavatus*) for recycling of KW into valuable compost. In this technology, vessel is made of nylon net with 90cm length and 35cm diameter containing 10-15 kg capacity plastic basket placed inside the vessel. This vessel can be hanged with the help of nylon rope in the available space of the house or on the branches of the tree planted around the house. KW were cut into small pieces (3-5 cm length) and kept inside the vessel followed by well decomposed cattle dung spread over the wastes. About 100-150 nos. of adult epigeic earthworms of three species in equal numbers is kept in the basket. Whole material is covered with jute bag and kept moist by adding water regularly to maintain the moisture. After 30 days the vermicompost gets ready for field application. The compost had organic carbon to total nitrogen ratio of 10-12: 1; 1.2% total nitrogen, 2.0% total phosphorus and 0.8% total potassium. This FNVC is an efficient technology and can be easily adopted to convert KW into valuable compost.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**


**A STUDY ON THE VARIATION OF THE PHYSICOCHEMICAL, HYDRO
BIOLOGICAL AND LIMNOLOGICAL PARAMETERS WITH RESPECTS TO
THE EUGLENA BLOOM AND ANTHROPOGENIC FACTORS IN A HIGH,
MODERATE AND LESS PRODUCTIVE POND OF BANKURA DISTRICT OF
WEST BENGAL, INDIA**

Subhajit Mahanty, Keshab Chakraborty, and Surajit Majumder*

Department of Zoology, Bankura Sammilani College, West Bengal, India

ABSTRACT

The Bankura town of district Bankura, West Bengal, India is full of lentic water bodies like pond, reservoir and water tank etc. Limnological characteristics of ponds in different seasons are related to the hydro biological condition of the Ponds. Physicochemical parameters of water and their influences on the occurrence of Euglena bloom and the distribution of other algal species. Sometimes water bodies are enriched with nutrients which lead to an excessive growth of plants that initially cause of Eutrophication and ultimately end up with algal bloom formation. Euglenophyte bloom is a common problem in most of the aquaculture ponds in Bankura. There is an intricate relationship between the metabolism of aquatic organisms and hydro-biological parameters in a fresh water body (Majumder and Dutta 2014). This study also focuses the impact of Euglena sp. on the growth and development of fish fauna. Algal bloom in the aquatic ecosystem is a natural process which triggered by different anthropogenic activity. Euglenophyte bloom is the common phenomenon in warmer shallow and eutrophic water bodies (Olaveson and Stokes, 1989; Xavier et al., 1991; Tripathi and Shukla, 1993) and it decreases the fish production and species diversity a lot. Our study is an attempt to study the dynamics of Euglena bloom in the various ponds of Bankura.



Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ASSESSMENT AND IDENTIFICATION OF EXOPHYTIC AND
ENDOPHYTIC BIOFLOC FORMING ORGANISM FROM SELECTED
MARINE ALGAE SOURCED ALONG THE COASTAL AREA OF
PANDURANGAPURAM, VISAKHAPATNAM**

**Arnepalli Chandrasekhar¹, C. Manjulatha¹,
Khadem Hussain Saedi¹ and D.V.V. Satynarayana Raju²**

Department of Zoology, Andhra University, Visakhapatnam, Andhra Pradesh

ABSTRACT

Nowadays, Biofloc technology is becoming both popular and successful in the field of aquaculture. Biofloc acts as a source of additional nutrients, improves immunity of the culture species and also maintains water quality by balancing the carbon and nitrogen ratio. In this context, there is urgent need to identify biofloc forming bacteria and therefore present study was undertaken. Initially marine algae belonging to Gracilaria sp., Ulva sp. and Cladophora sp. were selected to isolate exophytic and endophytic microbial population. For this, marine algal samples were collected from shallow waters of coastal area of Pandurangapuram, Visakhapatnam during the low tide period. Samples were sterilized and macerated and the solution obtained was subjected to serial dilution to isolate individual colonies. Isolated colonies were again subjected to quaternary streaking on Zobell Marine agar to get pure cultures. From these samples total 12 strains of exophytic and endophytic organisms were isolated aseptically, purified and preserved for further investigations. Preliminary identification of pure isolates was carried out by both culture and biochemical methods and was observed that majority of the organisms belong to genus Bacillus and Vibrio. Individual pure isolates were tested for their ability for floc formation under invitro conditions. In this experiment biofloc was formed with strain NPZ alone within two to three days of incubation. From harvested NPZ Biofloc, microbiological observation was carried out to know the purity of the floc formed. The whole experiment was repeated thrice and same results were obtained in every experimental setup. The NPZ strain was also able to form flocs in different salinity ranging from 1 ppt to 30 ppt. Preliminary Identification was carried out and strain NPZ was identified as Bacillus cereus based on culture and biochemical studies. By these findings, it was evident that bacterial strain NPZ belonging to the genus Bacillus had ability to form biofloc, which has potential application as inoculum in the field of Aquaculture and further studies are being carried out in both invitro and invivo conditions.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**BENTHIC FORAMINIFERA, SEDIMENT CHARACTERISTICS AND
ECOLOGICAL STUDIES IN THE NEAR SHORE ENVIRONMENTS OF PALK
STRAIT, TAMILNADU, EAST COAST OF INDIA**

M. Suresh Gandhi, K. Kasilingam, Annmaria K. George and Gayathri G.S.

Department of Geology, University of Madras, Chennai

ABSTRACT

A total of 72 offshore sediment samples in the depth range of 1-7m at Palk Strait, off Mandapam to Kodiyakarai were studied for Micropalaeontological investigations. This study highlights the presence of a composite cosmopolitan fauna of foraminifera of 112 benthic species belonging to 27 families and 57 genera from the 72 sediment samples. Among them *Rotalina* (49%) occupies a dominants followed by *Miliolina* (39%). The following genus *Ammonia*, *Asterorotalia*, *Quinqueloculina* dominates the living assemblages, followed by *Pararotalia* and *Elphidium*. From the total number of species, *Ammonia beccarii*, *Asterorotalia trispinosa*, *Pararotalia nipponica*, *A.tepida*, *A.dentata*, *Elphidium crispum*, and *Quinqueloculina seminulum* record profile abundance. However, only thirteen species namely *P.nipponica*, *Neorotalia calcar*, *Q.lamarckiana*, *Q.seminulum*, *Q.tropicalis*, *Triloculina trigonula*, *E.macellum*, *A.beccarii*, *E.crispum*, *A.dentata*, *A.tepida*, *Spirolloculina depressa*, and *S.communis* keep up a uniform distribution in more than six regions.

The influence of ecological parameters on the foraminifera taxa has been established in such a way that organic matter is contributed primarily by living species. The lack of relationship between the dead species and the organic matter suggests that the organic matter recorded in these sediments may have originated either from living benthic species or from plant remains reported in this Strait. Lower concentration of organic matter in the channel zones of 5-7m depth leads to disturbances of the seabed bottom due to the turbidity current movement. Significant concentrations of nearshore carbonates are mainly attributed to the deposition of transported carbonates mainly of broken shells and carried by the drifting currents.

From the present study, it is inferred that sediments are normally sandy in the Palk Strait but silty sand in deeper depths. All these factors help to conclude the progressive deposition or dispersion of sediments in seaward direction. The sandy nature of the Palk Strait regions outlines the deposition of sediments primarily by northerly moving currents. The presence of negative skewness along with the poor sorting confirms the existence of high energetic conditions in the depositional basins except for the northern and southern extremities. The alternate deteriorating sorting characters of the sediments in the different stations indicate the unevenness of the bottom topography. Geochemical analyses for Fe, Mn, Cr, Co, Ni, Pb, Zn and Cd were carried out. Based on the geochemistry, Fe concentration is very high compared to the all the trace elements. In general the concentrations of trace elements are higher at shallower depths and lower at deeper depths.

The present study has illuminated the fast prograding nature of the shelf, as evidenced by the living/dead ratios indicative of higher rate of sedimentation in this Strait. However, the lowest value of 7.50% at Manalmelkudi is attributed to a zone of low sedimentation due to its location on the limb of the spit which acts as a barrier to divert the influencing littoral currents.

As a highlight of the present study, it is concluded that more sediment deposition takes place in the southern part of the region than in the northern part of the sand bar where the total number of living species is much low comparatively other localities. The morphological deformative species also observed both northern and southern part of the study area. The trace elements concentration was much higher in the northern part of Manalmelkudi than in the southern part of Manalmelkudi, and it was concluded that the trace elements concentration fully affected the living population in the northern part of Manalmelkudi, so present study also brings out the dire need of protecting the gradually shallowing nature of Palk Strait as confirmed by the foraminiferal and geochemical evidences.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**CONSEQUENCES OF CLIMATE CHANGE ON THE
ENDANGERED BATS OF WESTERN GHATS, INDIA**

Sreehari Raman^{1,2*}, Shameer T.T.³, Pooja Ushakumari⁴

¹Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden
Chinese Academy of Sciences, Menglun, Mengla, Yunnan Province, PR China

²Department of Wildlife Sciences, College of Forestry, Kerala Agricultural University
KAU P.O., Vellanikkara, Thrissur, Kerala, India

³Advanced Institute for Wildlife Conservation, Tamil Nadu Forest Department
Vandalur, Chennai, Tamil Nadu, India

⁴College of Veterinary and Animal Sciences, Kerala Veterinary and Animal Sciences University
Pookode, Wayanad, Kerala, India

ABSTRACT

Western Ghats is home to 63 species of bats, including threatened and endemic species. Salim Ali's Fruit Bat (*Latidens salimalii*) is a cave-dwelling fruit bat restricted to the montane evergreen and adjoining coffee/cardamom plantations of the southern Ghats region. Due to various anthropogenic stress, there is a significant decline in the population of *L. salimalii* and now categorised under the endangered category by the IUCN. Climate change is an emerging threat to overall biodiversity, especially to the species with narrow or restricted distribution. Hence a study was conducted to identify suitable habitats for *L. salimalii* and the impact of climate change on the distributional ranges using MaxEnt. We followed the standard bat sampling protocol for collecting primary occurrence data. We also used secondary data to improve the accuracy and reliability of the models. Eco-geographic variables were identified based on the ecology of the species, and we used only the non-correlated variables for running final model. The habitat suitability model for *L. salimalii* was developed for current climatic conditions and projected their distribution for three Representation Concentration Pathway (RCP 4.5, 6.0, and 8.5) climate scenarios of the 2070 period. The results show that an overall 9,531 sq.km is suitable for *L. salimalii* under current climatic conditions. Variables like precipitation of the driest month, tree density, canopy height, and altitude have contributed much to the models. Future scenarios show a drastic decline in the highly suitable habitats across its distributional ranges. Our results thus inform managers to make a fitting move by looking at the appropriate example of this species to climatic change.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**CONSERVATION STRATEGIES OF RARE ETHNOMEDICINALLY
IMPORTANT ISODON NILGHERRICUS BENTH H. HARA PLANT**

Godse N.H. and B.M. Gaykar

Department of Botany, Ahmednagar College, Ahmednagar, Maharashtra

ABSTRACT

The present study focussed on ethnomedicinally important *Isodon nilgherricus* (Benth) H. Hara Basionym - *Plectranthus nilgherricus* Benth. *Plectranthus* species is well known for its medicinal and economic value. Studies revealed that it is a rich source of essential oil and diterpenoids. *Isodon nilgherricus* is an aromatic plant native to Africa and India. In India it is mainly found in southern ghats, Nilgiri hills, Ooty in Tamil Nadu District. Plants are grown with the help of seeds and saplings were successfully grown in Ahmednagar area as well in table land Wai in Maharashtra. In Tamil Nadu, Kurumba tribes especially in Nilgiri region the leaf paste of this plant is used against gum disorder and toothache. This study provides an initial information of conservation practices of *Plectranthus* species in Maharashtra.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**DEATH BY NETS: MONITORING BIRD
KILLS IN POMOGRANATE ORCHARDS**

Karthik N.J.¹, Vijaya Kumara² and Apoorva Kulkarni^{3,4}

Department of P.G. Studies and Research in Wildlife and Management
Kuvempu University, Jnana Sahyadri, Shankaraghatta, Shimoga, Karnataka

ABSTRACT

Fruit producers have identified bird damage as a critical issue as it incurs high costs. They try to limit this damage by deploying measures through traditional means or by using bird scaring techniques, reflecting devices, pesticides, and nets. Limited research has been carried out in monitoring the numbers and fate of the birds that get trapped in nets placed around fruit orchards. The present study was carried out with the objective of monitoring birds being caught in fishing nets used as a barrier to control the fruit damage by birds in pomegranate orchards in Chitradurga district of Karnataka, India. We assessed mortality rate of birds caught, noted their feeding guilds, and monitored them for six months from April to August 2021. A total of 36 species belonging to 11 orders and 20 families were recorded dead in the ten pomegranate plots studied. Some individuals were unidentified as they were decomposed with only few bones left in the nets. Among the 11 orders, Passeriformes were the most affected. A total of 1043 individual birds were observed entangled in the nets. Six individuals were found alive and were released after close observation of any injuries. Rose ringed parakeets were the most caught and found dead in the nets, followed by common myna and baya weavers. Feeding guilds of the birds entangled in nets comprised of majorly omnivores (44.4%) birds followed by insectivores (25%), granivores(16.6%), carnivores (11.11%) and frugivores (2.77%). Our study shows that monitoring mortality is essential in understanding populations of not only vulnerable and migratory species but also the common ones, which might not be common in the near future if such lethal bird control practices are not kept in check.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ECOSYSTEM SERVICES VALUATION OF A RAMSAR COASTAL LAKE
IN KERALA WITH SPECIAL EMPHASIS ON WATER QUALITY AND
LAND USE LAND COVER PATTERN**

Vishnu Sagar M.K, Sabu Joseph and Arunkumar P.S

Department of Environmental Sciences, University of Kerala
Thiruvananthapuram, Kerala, India

ABSTRACT

Coastal lakes and wetlands are dynamic ecosystems and provide many valuable [ecosystem services](#) (ESs) to human beings. The rapid increase in anthropogenic activities, socioeconomic development, and land use land cover (LULC) changes from the past two decades significantly affect the aquatic ecosystems and its ecosystem services. Unfortunately, these are not economically valued, especially in tropical and subtropical regions, which lead to under-investment and degradation of these ecosystems. This paper presents key results of a three-year international cooperation project focusing on ESs provided by Ashtamudi lake, a Ramsar site (Area= 56 km²), located in the coastal zone of Kerala, SW India. We present the results of water quality monitoring program in the lake and results of economic valuation of key ESs provided by the lake, including fisheries, tourism, transportation, mangrove vegetation. The economic valuation relies both on primary data collected on-site as well as data from secondary sources, like stakeholders workshops involving key representatives from the population, administration and academy and questionnaire based survey. The ESs based on fisheries, tourism, and transportation was estimated and is discussed here. The revenue generated from clam fishery (mainly *Paphia Malabarica*) and aquaculture activities are 6.099 million and 7.799 million USD respectively in 2017. The calculated value for carbon sequestration for mangroves (Area= 0.64 km²) is 144.64 Tonne C/year and revenue from this is 73314 USD in 2017, excluding flood management. The southern bank of the lake is densely populated and urbanized. The spatio-temporal variation of water quality attributes during non-monsoon and monsoon seasons show that DO, BOD and salinity exhibit spatio-temporal variations with adverse conditions on the seaward part of the lake. The southern zone shows hypoxic condition and Carlson Trophic State Index (CTSI) reveals that lake fell under eutrophic condition. The LULC around the lake basin were prepared and were classified into buildup area, agricultural land, wetlands, water bodies, and mangrove vegetation's and marsh lands. We found that all measured Land use classes and water quality characteristic correlations were statistically significant with the Ecosystem services; therefore, we concluded that Land use pattern and lake water quality affects the ecosystem services provided by the lake. These results could facilitate decision-making and development of related policies and might support finding strategies for sustainable management of the lake.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**INTERACTIONS BETWEEN INDIAN WILDLIFE AND
HUMANS IN TIGER STATE (MADHYA PRADESH)**

¹Yogesh Singh and ²Pawan Shijoria

¹Barkatullah University, Bhopal

²Govt. College Shahpur, Betul

ABSTRACT

Humans and wildlife interact "when animals offer a direct and repeated threat to the livelihood or safety of people, resulting in the persecution of that species," according to the World Wildlife Fund. In India, there is growing concern over such interactions. Today, there is a growing emphasis on human connection with wild animal both domestically and internationally. Both humans and wildlife are badly impacted, but many of the latter are endangered or threatened. With its ongoing research into not just the ecology of the wildlife but also how people respond to the presence of the animals in human-use settings, human interaction with wildlife has been at the forefront of this field of knowledge. India's scenario is, nevertheless, evolving. Wildlife habitats have been degraded and fragmented as a result of the growing human population and its need for natural resources. As a result, competition for the same few resources exists between people and wildlife. In addition to potentially undermining current and future conservation efforts, this change from coexistence to conflict could also make it more difficult to accomplish the Sustainable Development Goals (SDGs) and the Aichi Biodiversity Targets. The main challenges that are emerging in urban wildlife near protected areas are presented in the current study.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

NATURE-BASED SYSTEM FOR WASTEWATER TREATMENT AND REUSE

Monali Muduli

Academy of Scientific & Innovative Research (AcSIR) India
CSIR-Central Salt and Marine Chemical Research Institute, Bhavnagar, Gujarat, India

ABSTRACT

The water crisis is an alarming issue in the recent era. The population explosion, climate change, industrialization, and changing lifestyles of humans have created this global issue. Water stress varies significantly from place to place and can have a profound negative impact on public health, economic growth, and international trade. To meet the global demand for accessing pure water, it is necessary to follow sustainable approaches that manage the wastewater problem and help reuse the water. Traditional wastewater treatment systems' energy-intensive and mechanical components need significant upfront investment and high ongoing operational expenses. High maintenance costs and poor expertise in conventional treatment technologies have been unsuccessful in treating the wastewater adequately in developing countries like India. Compared with traditional approaches, nature-based technologies are pocket and user-friendly. Nature-based treatment system uses plants, soil, microbes, and other natural components to eliminate pollutants from wastewater. Thus, no chemicals and energy are used in the nature-based approaches, which makes it affordable, eco-friendly, and sustainable. Natural wetlands, constructed wetlands, and lagoons are nature-based systems used for various wastewater treatments. Water treatment and socio-economic benefits like increasing biodiversity, improving urban microclimates, control of flood and storm, biomass production, and water reuse could be achieved while treating water through nature-based systems. This paper will discuss the role of various nature-based systems in wastewater treatment and recent advancements developed in nature-based systems to enhance treatment efficiency.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**SPECTROPHOTOMETRIC MONITORING OF EXPLOSIVES WASTEWATER
TREATED USING INTEGRATED TREATMENT METHODS**

Bhumika Sharma^{1 2}, S. Mary Celin^{2*}, Garima Upreti² and Ranu Gadi¹

¹Indira Gandhi Delhi Technical University for Women, Kashmere Gate, Delhi

²Centre for Fire, Explosive and Environment Safety, (CFEES), DRDO, Delhi

ABSTRACT

Explosives are classified as potential human carcinogens as per USEPA. Activities such as manufacturing, transportation, destruction, loading assembly and packing of ammunition releases explosive chemicals into the environment. TNT red water majorly contributes to the mutagenic, carcinogenic and toxic properties of the effluent generated during TNT manufacturing. It contains high organic matter that interferes with its treatment methods. The main components of TNT red water are Dinitrotoluene sulfonate (DNTs), sodium nitrite, sodium sulfate, sodium sulfite and other nitro compounds contributing to its toxicity. Thus, there is a need to treat this wastewater for ensuring environmental safety. This study provides an integrated approach for the remediation of TNT red water which is divided into two steps; step 1 decolorization using sodium hypochlorite treatment followed by step 2 comparison of treatment methods viz. Zero valent ion, Fenton's oxidation integrated with advanced oxidation process. Monitoring of TNT wastewater treatment has been carried out by UV-Vis spectroscopy in terms of pH and its ionic concentration (nitrate and nitrite). Ionic concentration denotes the efficiency of the treatment process. Higher the ionic concentration better the degradation. Decolorizing TNT wastewater using liquid bleach followed by advanced oxidation process was found to be the most promising method with comparatively high nitrite ion concentration and complete treatment without the formation of precipitate or leaving any residue.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

WASTEWATER REUSE IN AGRICULTURE

Sumiran Dabrase¹ and Abhipsa Khobragade²

¹Department of Agriculture, R. B. College of Agriculture, Wardha (MH)

²Department of Agriculture, Rabindranath Tagore University, Bhopal (MP)

ABSTRACT

Water is an important natural resource all over the world. The environment, economic growth and development are all highly influenced by water. Scientists around the globe are working on new ways of conserving water. Rapidly depleting and elevating the level of freshwater demand, though wastewater reclamation or reuse is one of the most important necessities of the current scenario. Total water consumption worldwide for agriculture accounts 92% out of which about 70% of freshwater is used for irrigation which comes from the rivers and underground water sources. The statistics shows serious concern for the countries facing water crisis reported that 40% of the global population is situated in heavy water-stressed basins, which represents the water crisis for irrigation. It is an opportune time, to refocus on one of the ways to recycle water—through the reuse of wastewater, for irrigation in agriculture and other purposes.

Wastewater comprises of liquid wastes generated by households, industries, commercial sources, as a result of daily usage, production, and consumption activities. By giving primary and secondary treatments we can use this water for irrigation purpose in agriculture. There are various systems of wastewater treatments available. Wastewater and its nutrient content can be used extensively for irrigation and other ecosystem services. It's reuse can deliver positive benefits to the farming community. Therefore, wastewater reuse in agriculture is an ideal resource to replace freshwater use in agriculture. Treated wastewater irrigation supports agricultural yield and the livelihoods of millions of smallholder farmers. The rich availability of nutrients in reclaimed wastewater reduces the use of fertilizers, increases crop productivity, improves soil fertility.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**DEVELOPMENT OF REVERSE TRANSCRIPTION LOOP MEDIATED
ISOTHERMAL AMPLIFICATION (RT-LAMP) ASSAY AS A
POINT-OF-CARE DIAGNOSTICS FOR SARS-COV-2**

Shweta Kelkar¹ and Rashmi H.M.²

¹Dairy Microbiology Division, NDRI, Karnal, Haryana, India

ABSTRACT

The present study aimed to develop a reverse-transcription Loop-Mediated Isothermal Amplification (RT-LAMP) assay as a “Point-of-care” (PoC) diagnostic solution which can be useful for the detection of SARS CoV-2. To develop this assay, the complete genome sequences of SARS CoV-2 reported from different States and Union Territories of India were downloaded from GISAID (global initiative on sharing avian influenza data) database. The sequences were aligned to identify conserved regions for designing RT-LAMP primers. After the literature search, three conserved regions with no previously reported primers for detection of SARS CoV-2 were selected and the primers were designed uniquely using Primer Explorer V5. A total of six sets (A, B, C, D, E, and F) of primers, two each from the single conserved region were designed. The specificity of these primers was again checked *in silico* with BLAST tool in the NCBI server. The designed primers did not yield any similarity matches with the other human coronavirus except MERS-CoV and Bat SARS-CoV. On the other hand for the development of LAMP assay with these designed primers, the target conserved regions were got synthesized and analyzed for quality parameters before their use in the LAMP assay. Later, the primers (type, concentration, and ratio) and LAMP reaction temperature were optimized in the development of LAMP assay using synthetic gene constructs. Further, the performance of the developed assay was evaluated at KCGMC, Karnal, Haryana, using cDNA synthesized from RNA of SARS-CoV-2, and the developed LAMP technique was found effective in the detection of SARS-CoV-2. The developed LAMP technique was further explored for the development of colorimetric/fluorometric LAMP assay for rapid visual detection of SARS-CoV-2. The performance evaluation of developed colorimetric and fluorometric LAMP assays yielded satisfactory results in the detection of SARS-CoV-2 RNA. Hence, the developed Colorimetric and fluorometric LAMP assays can be used as PoC diagnostic tests for the detection of SARS-CoV-2. However, its performance directly with RNA or samples needs to be evaluated to make it an RT-LAMP assay and further validation is required with a larger sample size as per ICMR, India guidelines.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**LARGE SCALE SOIL MAPPING (1:10000) USING GEOSPATIAL
TECHNIQUES – A CASE STUDY OF DAHOD DISTRICT, GUJARAT**

**R. L. Meena^{1*}, P.C. Moharana¹, Mahaveer Nogiya¹, R. S. Meena²,
S. K. Yadav³, L. C. Malav¹, Brijesh Yadav¹, R. P. Sharma¹ and B. L. Mina¹**

¹ICAR-National Bureau of Soil Survey and Land Use Planning
Regional Centre, Udaipur, 313001, Rajasthan, India

²ICAR-National Bureau of Soil Survey and Land Use Planning, Regional Centre
Jorhat, 785004, Assam, India

³ICAR-National Bureau of Soil Survey and Land Use Planning
Regional Centre, New Delhi, 110012, India

ABSTRACT

The present study aimed to develop a reverse-transcription Loop-Mediated Isothermal Amplification (RT-LAMP) assay as a “Point-of-care” (PoC) diagnostic solution which can be useful for the detection of SARS CoV-2. To develop this assay, the complete genome sequences of SARS CoV-2 reported from different States and Union Territories of India were downloaded from GISAID (global initiative on sharing avian influenza data) database. The sequences were aligned to identify conserved regions for designing RT-LAMP primers. After the literature search, three conserved regions with no previously reported primers for detection of SARS CoV-2 were selected and the primers were designed uniquely using Primer Explorer V5. A total of six sets (A, B, C, D, E, and F) of primers, two each from the single conserved region were designed. The specificity of these primers was again checked *in silico* with BLAST tool in the NCBI server. The designed primers did not yield any similarity matches with the other human coronavirus except MERS-CoV and Bat SARS-CoV. On the other hand for the development of LAMP assay with these designed primers, the target conserved regions were got synthesized and analyzed for quality parameters before their use in the LAMP assay. Later, the primers (type, concentration, and ratio) and LAMP reaction temperature were optimized in the development of LAMP assay using synthetic gene constructs. Further, the performance of the developed assay was evaluated at KCGMC, Karnal, Haryana, using cDNA synthesized from RNA of SARS-CoV-2, and the developed LAMP technique was found effective in the detection of SARS-CoV-2. The developed LAMP technique was further explored for the development of colorimetric/fluorometric LAMP assay for rapid visual detection of SARS-CoV-2. The performance evaluation of developed colorimetric and fluorometric LAMP assays yielded satisfactory results in the detection of SARS-CoV-2 RNA. Hence, the developed Colorimetric and fluorometric LAMP assays can be used as PoC diagnostic tests for the detection of SARS-CoV-2. However, its performance directly with RNA or samples needs to be evaluated to make it an RT-LAMP assay and further validation is required with a larger sample size as per ICMR, India guidelines.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**BIODIVERSITY STUDIES OF PIGMENTED BACTERIA AND
FUNGI ISOLATED FROM PANJIM NALLAH WATER AND
CHARACTERISATION OF SELECTED ISOLATES**

Marielou Ferrao

Assistant Professor Department of Microbiology
St. Xavier's College of Arts Science & Commerce Mapusa Goa

ABSTRACT

The St. Inez nallah is an aquatic freshwater ecosystem that originates from the Altinho hill at Taleigao, branching along the way through agricultural lands and the city market, opening up in the river Mandovi. The water is contaminated by a huge amount of sewage which enters the nallah from the illegal community residences along the nallah. In this scenario, a biodiversity study has crucial value. Previous studies have shown that pigmented microorganisms have been isolated from polluted aquatic environments; hence a biodiversity study of pigmented bacteria and fungi in this ecosystem becomes extremely relevant. Isolation of pigmented bacteria and fungi was achieved by membrane filtration of collected water samples on Kings Medium and Potato Dextrose agar with glycerol. Pigmented isolates were screened for biosurfactant production by cetyltrimethylammonium bromide plate methylene blue assay. Isolates showing potential for accumulation of biosurfactant were chosen for determination of taxonomic status by 16SrRNA sequencing. Optimisation of growth in mineral medium was achieved with glycerol as a carbon source. Out of the 14 pure cultures of pigmented bacteria and fungi obtained two isolates SXC3 and SXC31 showed potential for production of biosurfactant. The culture SXC31 gave maximum growth at glycerol concentration of 0.02% and culture SXC3 gave maximum growth at glycerol concentration of 0.5% when glycerol was the sole carbon source in mineral medium. Cultures SXC3 and SXC31 did not display pigmentation when glycerol was the sole carbon source in mineral medium. SXC3 was identified as *Klebsiella michiganensis* strain BAE24 and SXC31 was identified as *Pseudomonas entomophila* strain W-S-1-1-2 by 16SrRNA sequencing.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



Technical Session

4



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ASSOCIATION OF ABO BLOOD GROUP PHENOTYPES WITH
COVID-19: A CROSS-SECTIONAL STUDY IN GOA**

Neeha Sinai Borker and Nandini Vaz Fernandes

Department of Zoology
Parvatibai Chowgule College of Arts & Science (Autonomous), Gogol, Goa, India

ABSTRACT

COVID-19 pandemic has caused massive havoc across millions of people worldwide. At present, extensive research is being emphasized on vaccine trials, treatment, and risk factors for the disease. Recent studies suggest one such inheritable trait, ABO blood group phenotype, may be associated with SARS-CoV-2 infection. The present study is aimed at determining the susceptibility and symptomatic variation of COVID-19 among ABO blood group phenotypes in a sample population of Goa, India. An online questionnaire-based survey was conducted among 228 Goan subjects by administering a google form. Subjects who were tested positive for COVID-19 and had information about their blood groups were recruited for this study. The data collected was subjected to statistical analyses using Fisher's exact test. Present study showed subjects with Blood group O were at a higher risk for COVID-19 (OR= 1.4 ,95% CI, 0.93-2.22, p-value=0.114), followed by B (OR= 1.4 ,95% CI, 0.93-2.22, p-value=0.896), A (OR= 1.4 ,95% CI, 0.93-2.22, p-value=0.488), while AB was at lower risk (OR= 0.43 , 95% CI, 0.17-1.13, p-value=0.090). Additionally, Rh+ subjects were at a lower risk for COVID-19 (OR= 0.77, 95% CI, 0.37-1.68) as compared to Rh- subjects (OR= 1.29, 95% CI, 0.59-2.64). However, the difference observed was statistically insignificant (p=0.460). Our findings suggest that ABO blood group phenotypes may influence the susceptibility of people towards COVID-19.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**ENDOPHYTIC FUNGI INOCULATION: A SUSTAINABLE WAY
FOR INCREASING THE PHYTOCHEMICAL CONTENT IN
STEVIA REBAUDIANA (BERTONI) BERTONI**

Rachna Verma and Reema Devi, Kiran

School of Biological and Environmental Sciences
Shoolini University of Biotechnology and Management Sciences, Solan, Himachal Pradesh, India

ABSTRACT

The demand for *Stevia rebaudiana* has increased due to its low calorie and medicinal values. *S. rebaudiana* a perennial herb belonging to family asteraceae is important for its leaves which are used as non-calorie sweetener and in many therapeutic applications including diabetes. Endophytes have the influence on biological properties of medicinal plants, so the research is focused on twenty fungal isolates classified as ascomycota (R1-R14), zygomycota (R15-R17), or oomycota (R18-R20) which were isolated from the medicinal plant *Ocimum tenuiflorum* L. Among all fungal isolates R2 (*Fusarium fujikuroi*) strain showed maximum antagonistic activity towards plant pathogenic fungi *Rosellinia necatrix* and *Fusarium oxysporum*. Fungal endophyte *Fusarium fujikuroi* (ON652311) was inoculated on *Stevia rebaudiana* seeds to determine the effect of an endophytic fungus on the phytochemical properties of *Stevia rebaudiana*. During the study, it was observed that the highest amount of total phenolic, flavonoids, tannin, carbohydrate, protein alkaloid, saponin compounds, were recorded from the methanolic extract of endophyte inoculated plant. The lowest amount of total phenolic flavonoids, protein, tannin, carbohydrate, alkaloid, saponin, compounds were recorded in chloroform extract. GC-MS analysis of methanol plant extract of *S. rebaudiana* plant showed eleven prominent peaks. The major compound observed in methanol plant extract were Styrene (C₈H₈) with peak area 13.62%, 1,3-Benzodioxole-2-carboxylic acid, ethyl ester (C₁₀H₁₀O₄) with peak area 13.25. In chloroform plant extract, the major compound observed were 1H-Imidazole-1-ethanol, à,á-diphenyl-, (R*,R*)- (C₁₇H₁₆N₂O) with peak area 8.20%, N-(3-Phenylbutyl)-pentadecafluor o-octanamide (C₁₈H₁₄F₁₅NO) with peak area 6.82%. Thus, present research proved that methanol acts as best solvent to extract maximum quantity of phytochemicals from the plant and as well proved endophytic fungi inoculation as a sustainable way for increasing the phytochemicals in plants.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**SENSITIVITY OF *ANDROGRAPHIS PANICULATA* (BURM. F.) MORPHOTYPES
TO SOIL MOISTURE STRESS: STATUS OF ANTIOXIDANT ENZYME
ACTIVITIES AND ANDROGRAPHOLIDE YIELD**

**Kuldeep Singh A. Kalariya, Deep Shahia, Parmeshwar Lal Sarana,
Ram Prasanna Meena¹, Narendra Gajbhiyea, Rohan Sarkar¹, and Geetha K.A.¹**

ICAR-Directorate of Medicinal and Aromatic Plants Research, Boriavi, Gujarat

ABSTRACT

Well known across the world for its use in treatment of several diseases and alignments, the *Andrographis paniculata* (Burm. f.) Wall is popular for its bitterness. Steroidal lactones; andrographolide, neoandrographolide and dehydroandrographolide are major bioactive compounds from the entire herbage of this plants. As per reports, it is sensitive to water deficit stress and hence, with a view to test stress mitigating effects of foliar exogenous application of regulatory phytohormones; methyl jasmonate (MeJ) and salicylic acid (SA), five different morphotypes were evaluated under soil moisture stress conditions imposed during 90-140 days after transplanting in two consecutive crop seasons. The leaf water content (RWC) content and leaf water potential (LWP) were not affected by MeJ and SA however, soil moisture stress had severely affected the soil water plant relations in all morphotypes. Stress had significantly upregulated the activities of activity of superoxide dismutase (SOD), guaiacol peroxidase (GPOX), catalase (CAT) and ascorbate peroxidase (APX). Stressed plants had compromised dry herbage yield but, increase in andrographolide concentration by 6% however, the net loss in terms of andrographolide yield was nearly 8% in stressed plants. Foliar exogenous application of MeJ and SA mitigated stress effects through enhanced antioxidant enzymes activity, however, with no significant effects dry herbage yield in any morphotypes.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**A NEW CALIX[4]PYRROLE FUNCTIONALIZED FLUORESCEIN BASED
SUPRAMOLECULAR COMPOUNDS FOR LIQUID CRYSTALLINE AND
WINDOW LAYER SOLAR CELL APPLICATIONS**


Suryajit L. Rathod¹, Hitesh M. Parekh^{1,2*} and Vinay S. Sharma^{1*}

¹Department of Chemistry, School of Science, Gujarat University, Gujarat, India

²Department of Chemistry, Sardar Patel University, Vallabh Vidhyanagar, Gujarat, India

ABSTRACT

A newly symmetrical liquid crystals (CPFL₁-CPFL₄) based on calix[4]pyrrole as a central rigid core were synthesized *via* esterification reaction. All four functionalized compounds exhibits columnar hexagonal phase over a higher mesophase range and further stabilized upto room temperature. The thermal behaviours and optical textures were identified by using DSC, POM and the molecular organization of the compound in the mesogenic state is confirmed by XRD technique. Calix[4]pyrrole core with symmetrical nature exhibited columnar type self-assembly at room temperature. All four supramolecules with different side spacer showed higher thermal stability. Compound CPFL₁ is further tested for the absorbing layer in solar cell applications specially replace toxic CdS layer in solar cell devices. Herein, we used calix[4]pyrrole functionalized supramolecular liquid crystalline compounds as an optical window layer to achieve higher transmittance behavior towards the incident irradiance to the absorber layer. The current-voltage relation shows Ohmic-type behaviour. The photophysical behavior display high transmittance in visible and nearly IR regions. The derivative showed lower band gap values at different thin film state and variable temperature which are more suitable for ecofriendly absorbing window layer in solar cell applications. The morphology clearly indicates the uniform deposition of the sample with grain-growth type arrangement pattern.



Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**EFFECT OF FENUGREEK SEED EXTRACT ON FLUORIDE
INDUCED TESTICULAR DISORDERS IN ALBINO RATS**

Rajkumar Maiti*¹ and Debidas Ghosh²

¹Department of Physiology, Bankura Christian College, Bankura, West Bengal, India

²Department of Bio-Medical Laboratory Science and Management
Vidyasagar University, Midnapore, West Bengal, India.

ABSTRACT

Fluoride is one of the most abundant trace elements present in nature especially in rocks, soil, coal and drinking water. Fluoride contamination in drinking water have been documented as one of the greatest human health issues globally. According to WHO and BIS fluoride concentration up to 1-1.5 mg/L is recommended as a permissible limit for drinking. Prolonged intake of fluoride through drinking water above this limit is associated with dental and skeletal Fluorosis. Fluoride contamination through drinking water can disrupt male gametogenesis and steroidogenesis and also induce testicular oxidative stress. The aim of this study was to investigate the efficacy of fenugreek seeds as a potential natural source of antioxidants. A fenugreek seed (*Trigonella foenum-graecum L.*) is a member of the leguminous family and its seeds are used as herbal medicine. This study was carried out to evaluate the effect of aqueous extract of fenugreek seeds at the dose of 80mg/100g body weight/day for 28 days on fluoride-induced testicular toxicity in albino rats. Treatment of rats with sodium fluoride at the dose of 2 mg/100g/day for 28 days resulted in a significant reduction of testicular ⁵, 3-hydroxysteroid dehydrogenase and 17-hydroxysteroid dehydrogenase activities along with low plasma levels of testosterone, follicular stimulating hormone and luteinizing hormone. On contrary, fluoride treatment was associated with diminished activities of testicular catalase, superoxide dismutase and peroxidase along with elevation of thiobarbituric acid reactive substances and conjugated dienes. Co-administration of fenugreek seed to the fluoride treated animals resulted in a significant recovery from testicular disorders. In addition, fenugreek seed treatment led to a remarkable decrease in the level of lipid peroxidation and increase in the activities of antioxidant enzymes. Thus, it can be concluded that fenugreek seed extract can protect the sodium fluoride-induced testicular toxicity and this effect may be attributed to its antioxidant properties.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**SEMECARPUS ANACARDIUM LINN. LEAF EXTRACT EXHIBITS
ANTI-BREAST CANCER PROPERTIES AND PROLONG THE
LIFE-SPAN OF TUMOR-BEARING MICE**

Rajesh Kumar Singh^{1*}, Bhagaban Mallik², Amit Ranjan¹ and Anil Kumar Singh¹

¹Department of Dravyaguna, Institute of Medical Sciences, Banaras Hindu University, Varanasi, India

²Department of Anatomy and Cell Biology, University of Iowa, Iowa City, USA

³Department of Biochemistry, Institute of Science, Banaras Hindu University, Varanasi, India

ABSTRACT

Semecarpus anacardium Linn. nuts are used in preparation of drugs of several traditional medicine system to treat clinical ailments since ancient times that adversely affect the population of this plant, and finally the biodiversity. The alternative source for the nuts is demand of the time to preserve the biodiversity without affecting the clinical demands.

The phytochemical components of nuts can be difficult to isolate and have lethal effects on other cells. We used standardised techniques in this study to isolate the phytochemicals from the leaf extract and assessed their anti-cancer properties for potential drug development. The ethyl acetate leaf extract selectively affects cancer cells in a dose-dependent manner (IC₅₀: 0.57 µg/ml in MCF-7 cells) in various cancer cell lines. However, the non-malignant cells were relatively insensitive to the extract. Further study with acridine orange, ethidium bromide and 4',6-diamidino-2-phenylindole (DAPI) revealed that the extract induced apoptosis in cancer cells.

Next, we examined if the extract incubation could induce cell cycle arrest and suppress cell migration in the cell culture model. Consistent with this idea, the leaf extract could potentially affect the aggressive nature of migration in cancer cells. Moreover, oral administration of extract significantly restored the tumor growth of Ehrlich ascites carcinoma cells in mice. Besides, the ethyl acetate extract could potentially prolong the survival of tumor-bearing mice.

Together, these observations suggest the anti-cancer activities of *S. anacardium* leaf potential for both *in vitro* and *in vivo* models. Hence, this study opens a new paradigm to examine the phytochemical constituents of the leaf for anti-cancer activities.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**EVALUATION OF NEUROPHARMACOLOGICAL ACTIVITIES OF
PERSIMMON (*DIOSPYROS KAKI* LINN.) FRUITS IN RATS**

Asmita Arondekar, Sweaton Fernandes, Nevita D. Rivonkar, Mithila R. Mhamal

Goa College of Pharmacy, 18th June road, Panaji-Goa

ABSTRACT

The *Diospyros kaki* L., also known as oriental persimmon, is a tropical, deciduous fruit with a fleshy, fibrous texture that is a member of the Ebenaceae family. The fruit is rich in polyphenols, terpenoids, steroids, flavonoids, carotenoids, minerals, dietary fiber and other bioactive compounds.

The objective of the study is to conduct pharmacological screening of the persimmon (*Diospyros kaki* L.) aqueous and ethanolic extract in order to assess the extract's impact on the rat CNS utilizing various animal models.

The AEDK and EEDK were screened for antianxiety activity, spontaneous behavior and effects on learning and memory using different animal models. Two doses each of AEDK (200mg/kg and 400g/kg) and EEDK (200mg/kg and 400mg/kg) were selected for the study.

From the data, it can be inferred that the test groups have antianxiety activity because all four test groups increased in metrics such %OAE and %TSOE at some point in time. AEDK had more antianxiety effect than the fruit's ethanolic extract. In conclusion, pharmacological testing revealed that the aqueous and ethanolic extract of *Diospyros kaki* L. significantly reduced anxiety and had CNS depressive effects.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

HYPERTENSION: A MAJOR THREAT TO HUMAN BEINGS

Supriya Meena* and Jaya Arora

Laboratory of Biomolecular Technology, Department of Botany,
Mohanlal Sukhadia University, Udaipur, Rajasthan (India)

ABSTRACT

Nowadays especially after Covid-19 pandemic, stress full lifestyle has led to the high blood pressure conditions which in turn resulted in hypertension. Almost half of all adults in the United States have high blood pressure, but many are not aware of this fact. Hypertension is a primary risk factor for cardiovascular disease, including stroke, heart attack and heart failure. Keeping blood pressure under control is vital for preserving health and reducing the risk of these dangerous conditions. The U.S. National Heart, Lung, and Blood Institute (NHLBI) recommend the DASH diet for people with high blood pressure. DASH stands for “Dietary Approaches to Stop Hypertension.” Angiotensin-I-converting enzyme (ACE; peptidyl dipeptide hydrolase, EC 3.4.15.1) has a key role in the control of blood pressure through the renin angiotensin-aldosterone system. The key role of ACE is to convert angiotensin-I (the inactive decapeptide) to angiotensin-II (the potent vasoconstrictor octapeptide). ACE-inhibitory activities of peptides can be affected by their chain length, composition and sequence like conformational characteristics. ACE-inhibitory peptides have their potential of stability in gastrointestinal tract and reach the digestive, endocrine and cardiovascular system to show their bioactivity. Administration of antihypertensive medicines such as captopril and enalapril, which are potent ACE inhibitors, are among the synthetic medicines have shown to have undesirable side effects such as insomnia, angioedema, cough, allergic reactions, fever and headache. Antihypertensive peptides/protein hydrolysates have been reported as natural ACE inhibitors that are also equally capable of reducing blood pressure with no side effects. Natural antihypertensive peptides have been reported from several medicinal plants such as *Benincasa hispida* and *Glycine max*. With that view many of plant bioactive peptides will become potent sources of such natural antihypertensive protein hydrolysates and help us to disease risk reduction.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**HYPO-TESTICULAR ACTIVITY ASSESSMENT OF HYDRO-METHANOL
(60:40) SEED EXTRACT OF *LUFFA ACUTANGULA* IN HUMAN AND
ALBINO RAT: AN *IN-VITRO* INVESTIGATION**

Tanusree Mondal* and Debidas Ghosh

*Nutrigenomic Research Laboratory, Dept. of Bio-Medical Laboratory Science & Management
Vidyasagar University, West Midnapore, West Bengal, India

ABSTRACT

Unprecedented population growth is one of world's major issues and it is most important priority area of research to control population growth. This *in-vitro* experiment was designed to explore the male contraceptive efficacy in different dose (10mg, 20mg and 40mg) of hydro-methanolic (60:40) extract (HME) of seed of *Luffa acutangula* in rat and human reproductive parameters. Spermiological sensors i.e., sperm motility, viability, HOS, acrosomal cap status were assessed in rat and human spermatozoa and also nuclear chromatin decondensation (NCD) were assessed in rat spermatozoa. Activities of oxidative stress biomarkers i.e., SOD, catalase and free radical end product i.e., TBARS also measure from testicular and epididymal tissue of rat and human sperm pellet. Activities of testicular androgenic key enzymes i.e., $\Delta 5,3\beta$ -HSD and 17β -HSD were assessed in rat. Toxicity profile i.e., GOT and GPT were assessed from various tissue samples in rat like liver, kidney, testis and prostate. All spermiological profile were reduced significantly ($p < 0.05$) with different doses (10mg, 20mg, 40mg). EC50 value determines that fifty percent immobilization in spermatozoa as it was noted in human at 10mg/0.5ml dose and in case of rat the EC50 was 8mg/0.5ml dose of seed extract. The testicular androgenic key enzymes activity i.e., $\Delta 5,3\beta$ -HSD, 17β -HSD were reduced significantly ($p < 0.05$). The activities of SOD and catalase were significantly ($p < 0.05$) reduced and also significantly ($p < 0.05$) increased the level of TBARS in rat testicular and epididymal tissue and human sperm pellet. No metabolic toxicity was noted on different reproductive and metabolic organs of rat. *In-vitro* investigation concludes that hydro-methanol (60:40) seed extract of *L. acutangula* possess a significant anti-testicular cum contraceptive efficacies in human and rat.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**INTERACTIVE EFFECT OF ELEVATED CO₂ AND VARYING
NITROGEN REGIME ON THE PHYSIO-BIOCHEMICAL
RESPONSE OF *NEOLAMARCKIA CADAMBA***

Manish Singh¹ and Hukum Singh²

Department of Forest Ecology and Climate Change
Forest Research Institute, Dehradun

ABSTRACT

Quantitative integration of the literature on the effect of increasing CO₂ and nitrogen regimes on woody plants is crucial to aid our understanding of forest health in the future decades and to better forecast terrestrial feedbacks on the global carbon cycle. *Neolamarckia cadamba* is one of the promising, multipurpose, prospective and fast growing tree which is being grown in field in form of agroforestry. *N. cadamba* is considered to be a highly valuable tree in agroforestry and Carbon Sequestration due to its quick decomposition rate. However, there is no information on how this plant will react to changing climatic circumstances, including higher atmospheric CO₂ concentrations. The aim of the study is to understand the physio-biochemical response of *Neolamarckia cadamba* under elevated carbon dioxide (~800 μmol mol⁻¹) as compared to control (400 μmol mol⁻¹) with three different nitrogen regimes. Protein were increased significantly in N₃₀₀ Kg N ha⁻¹ in elevated CO₂ concentration while chlorophyll and carbohydrate were significantly increase in N₅₀₀ Kg N ha⁻¹ in elevated concentration. While plant respiration, were increased significantly in N₂₀₀ Kg N ha⁻¹ in elevated CO₂ concentration as compared to control. Plants maintain stronger stress tolerance mechanism via improvements in the synthesis of photosynthate and water retention capacity. Such behavior may facilitate plant adaptation to deal with climatic change and global warming. Tree productivity, vegetation, and ecosystem health may also be improved.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**NOVEL MECHANISTIC INSIGHTS ON MELATONIN MEDIATED PHYSIO-
BIOCHEMICAL, MICROSCOPIC, AND HISTOCHEMICAL MODIFICATIONS
IN TOMATO PLANTS TO PROMOTE SALT STRESS TOLERANCE**

Tanveer Ahmad Khan¹ and Qazi Fariduddin^{1*}

¹Plant Physiology and Biochemistry Section, Department of Botany
Faculty of Life Sciences, Aligarh Muslim University, Aligarh, India

ABSTRACT

Salt stress is a serious environmental constraint that is increasingly limiting seed germination, plant growth and development as well as both yield and quality of agricultural crop plants. Melatonin (Mel) is an indolamine multi-functional signaling molecule governing diverse range of environmental stresses responses in plants. Therefore, an experiment was designed to explore the defensive role of Mel under salt stress conditions. Root dip pretreatment of Mel (50 or 100 μ M, 4h) was applied to tomato plants just before transplantation and NaCl (100 or 150 mM) stress was imposed at 20 days after transplantation. It was observed that salt stress adversely affected stomatal physiology, photosynthetic efficacy, plant growth, and disrupted ultra-structure of chloroplast, root morphology, and cell viability, however, salt stress triggered hyper-accumulation of reactive oxygen species (H_2O_2 and O_2), and negatively affected membrane integrity. Interestingly, pretreatment of Mel significantly reduced the MDA content and electrolyte leakage, mitigated oxidative stress and promoted photosynthesis, improved cell viability, stomatal behavior, and plant growth, and boosted the activity of catalase (CAT), peroxidase (POX), and superoxide dismutase (SOD) and elevated proline content which bestowed endurance and alleviated the salt-induced toxicities.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**SEQUENCE ANALYSIS OF *siet* gene IN *STAPHYLOCOCCUS
PSEUDINTERMEDIUS* ISOLATED FROM CANINE PYODERMA**

**Prashantha M K; Shambulingappa B E*; Sundareshan S;
Kotresh A M; Rudresh B H; Madhavaprasad C B and Arun S J**

Veterinary college Shimoga (KVAFSU, Bidar), Karnataka

ABSTRACT

A study was conducted on isolation, identification and characterization of staphylococcal organisms associated with the canine pyoderma. The bacteriological processing of the samples resulted in the recovery of 95 staphylococcal isolates and 18 other bacterial isolates. On culture, staphylococci were the most predominantly (n=95, 75.39%) isolated organisms. Based on *nuc* gene-based PCR, out of 95 staphylococcal isolates obtained, 82 (86.1%) of the isolates were found belonging to *S. pseudintermedius*. And out of 82 *S. pseudintermedius* isolates, *siet* gene was detected in 69 (86.1%) isolates. *S. pseudintermedius* was found to be predominant bacterial pathogen responsible for pyoderma in dogs.

Two PCR products which were amplified for *nuc* gene and one each of *siet* and *mecA* gene were sent for commercial sequencing and the sequence alignment revealed 99.23 per cent, 99.14 per cent, 98.98 per cent and 100 per cent, similarities with the corresponding genes of *S. pseudintermedius* respectively.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**SIMULTANEOUS ESTIMATION OF METHOCARBAMOL AND IBUPROFEN
IN FIXED DOSE COMBINATION BY UV SPECTROPHOTOMETRIC METHOD**

Premanand Velip¹, Anand Mahajan¹, Adison Fernandes¹ and Shweta Borkar^{1*}

¹Department of Pharmaceutical Analysis
Goa College of Pharmacy, Panaj, Goa, India

ABSTRACT

The objective of present research work is to develop a simple, precise and accurate UV spectrophotometric method based on Isobestic point approach for quantitative estimation of methocarbamol (MET) and ibuprofen (IBU) in their fixed dose combinations. All solutions were prepared using methanol as solvent. The isobestic point wavelength for MET and IBU was found to be 226 nm. Concentration of IBU was determined at isobestic wavelength 226 nm while MET was determined directly at 277 nm as there was no interference of IBU. Validation of the method was carried out as per ICH guidelines (Q2R1). Linearity was established for MET at the concentration range of 28 – 56 µg/mL ($r^2 = 0.996$ at 277 nm) and for IBU it was between 16 – 32 µg/mL ($r^2 = 0.990$ at 226 nm). Percentage relative standard deviation (% RSD) value of less than 2 % for intra-day and inter-day precision confirmed the precision of the method. Accuracy of the method was in the range of 98 - 102 %. The developed method is suitable and can be applied for analysis of marketed formulation.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**STUDIES ON CLINICAL SPECTRUM OF CANINE PYODERMA ASSOCIATED
WITH *STAPHYLOCOCCUS PSEUDINTERMEDIUS* AND OTHER SECONDARY
BACTERIAL INVADERS AND THEIR ANTIBIOGRAM PROFILE
FOR EFFECTIVE THERAPY**

**Prashantha M K; Shambulingappa B E*; Sundareshan S;
Kotresh A M; Rudresh B H; Madhavaprasad C B and Arun S J**

Veterinary college Shimoga (KVAFSU, Bidar), Karnataka

ABSTRACT

A study on bacteriological investigation of canine pyoderma cases was conducted at the Veterinary College, Shivamogga. Exudate/pus/lesion swabs were collected from clinical cases of canine pyoderma (n=126) and subjected to isolation and identification of bacterial isolates by phenotypic methods. The bacteriological processing of the samples resulted in the recovery of 95 staphylococcal isolates and 18 other bacterial isolates. On culture, staphylococci were the most predominantly (n=95, 75.39%) isolated organisms. The isolates obtained other than staphylococcal species were *E. coli*, *Pseudomonas* species, *Klebsiella* species, *Proteus* species. Subsequently *in vitro* antimicrobial sensitivity testing was carried out for all the isolates. *S. pseudintermedius* showed highest susceptibility to clindamycin and the highest resistance to enrofloxacin, and the other bacterial agents were found sensitive to co-trimoxazole and ceftriaxone and resistant to amoxicillin/sulbactam and enrofloxacin. A study on the occurrence patterns of canine pyoderma revealed higher frequency of cases in male dogs, in the age group of 1-2 years and in Labrador breed. We report the *S. pseudintermedius* as the most predominant pathogen associated with the canine pyoderma in the study area.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**DEVELOPMENT OF LOW COST
PISTON PRESS BRIQUETTING MACHINE**

M. Z. Sheikh¹, A. G. Mohod² and Sakhi Katre³

Director
ICAR-National Research Centre on Camel, Bikaner, India

ABSTRACT

The two most-promising climate resilient species of camel thrive well in hot arid and semi-arid regions of the desert (e.g. *Camelus dromedaries*) and arid cold-climatic regions of the mountains (e.g. *C. bactrianus*) and continue to provide livelihood opportunity to co-habiting human population. They serve as multipurpose animals producing milk, fiber/hair, as a means of transport or draught, excreting urine and providing manure, often yielding meat as a part of culling process and supplying leather and bones after death. The unique ability to adapt to extreme desert ecosystem with peculiar physiological (thermoregulation, water metabolism, glucose and energy metabolism, salt tolerance, forbearance against choking dust, etc.) and anatomical (fore limb and hind limb, long neck, single and double hump, third eye-lid, forestomach, etc.) differences has a significant bearing on its productive lifespan. Methane emission from ruminants is a significant component of greenhouse gas (GHG) production that contributes to global warming, but camel seems to contribute less due to its unique fiber-utilizing capability and foraging on phytochemical-rich forages that controls methane emission. Thus, it is noteworthy to say that camel is an environmental-friendly livestock of the globe and its remarkable adaptive characteristic projects it as the animal for future as the world is preparing itself to face the untoward challenges of climate change.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**DEVELOPMENT OF BIODEGRADABLE NANOCOMPOSITE
PACKAGING MATERIAL FOR CHICKEN MEAT**

Kandeepan. G*., Monish. V., Sowndarya. A., Babji. Y., Spoorthi A., Srikanya., Ayesha. A

ICAR-National Research Centre on Meat, Hyderabad, Telangana-

ABSTRACT

Meat is a moisture-rich nutritious animal protein having great demand among consumers. The recent food packaging regulations amidst climate change recommend the usage of biodegradable packaging material for foods for preventing environmental pollution. Biopolymer-based films have a poor barrier and thermal properties compared to plastic films. This makes exclusive biopolymer films ineligible for use as primary packaging material for moisture-rich food products like meat. Hence research was undertaken to develop a biodegradable nanocomposite-biopolymer film for improved barrier and thermal properties for packaging meat. The eco-friendly nanocomposite packaging material was developed by incorporating polylactic acid, nano clay, and clove oil in chloroform. The film was prepared by casting method in a large glass tray. The nanocomposite film was characterized for quality parameters in comparison with the commercial low-density polyethylene (LDPE) films. The results indicated that film thickness, transparency, tensile strength, and lightness were significantly ($p < 0.05$) higher for nanocomposite film when compared with the LDPE film. Whereas, traits like elongation at break were significant ($p < 0.05$) in LDPE film compared to nanocomposite film. There was no significant difference in the parameters like moisture content, moisture absorption, water vapor permeability, and film solubility between the LDPE and nanocomposite films. Then the developed nanocomposite film was made into pouches to hold 250g of chicken meat. The pouches were sealed and the meat was stored in a refrigerator at $4 \pm 1^\circ\text{C}$ to analyze the shelf-life attributes of the chicken meat. A control group was maintained with chicken meat packaged in LDPE pouches. The storage quality attributes revealed that there was no significant difference between the chicken meat packaged in LDPE and nanocomposite films for the parameters like pH, titratable acidity, and thiobarbituric acid reactive substances. While the parameters like extract release volume and tyrosine values differed significantly ($p < 0.05$) between the chicken meat packaged in LDPE and nanocomposite films during the refrigerated storage period. The shelf-life attributes showed that chicken meat was acceptable upto day 6 in both LDPE and nanocomposite films. It is concluded from the above results of film characteristics and shelf-life attributes that the developed nanocomposite film is an ideal alternative for LDPE films for packaging fresh chicken meat and subsequent storage in refrigerated conditions.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**DEVELOPMENT OF ELECTROCHEMICAL BASED IMMUNOSENSOR
FOR THE DETECTION OF COLISTIN-A BANNED ANTIBIOTIC IN
CHICKEN MEAT**

Harsh Kumar¹, Kamil Kuča², and Dinesh Kumar^{1*}

¹School of Bioengineering and Food Technology
Shoolini University of Biotechnology and Management Sciences, Solan, India

²Department of Chemistry, Faculty of Science
University of Hradec Kralove Hradec Kralove, Czech Republic

ABSTRACT

An innovative amperometric immunosensor has been developed to detect antibiotic colistin from the chicken liver. Colistin is an antibacterial peptide that has been barred for human consumption, but it is being commonly used as a veterinary drug, and as a feed additive for livestock. In the present work, an immunosensor was developed by immobilizing an anti-colistin antibody onto the carbon nanofibers-gold nanoparticles surface of the screen-printed electrode. The sensor records electrochemical response in the chicken liver spiked with colistin with cyclic voltammetry (CV). Additionally, the characterization of electrode surface was done with Field-emission scanning electron microscope, Fourier transform infrared spectroscopy, and Electrochemical impedance spectroscopy at each step of fabrication. The lower limit of detection was $0.89 \mu\text{gKg}^{-1}$, with a R^2 of 0.901 using CV. Further validation of the immunosensor was conducted using commercial chicken liver samples, by comparing the results to those obtained using traditional methods. The fabricated immunosensor showed high specificity towards colistin, which remained stable for 6 months but with a 13% loss in the initial CV current.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**DPP-IV INHIBITION (ANTI-DIABETIC)
POTENTIAL OF KAPPA CASEIN HYDROLYSATES**

Harshita Sonarshi, Sathish Kumar M H and Rajani C S

Southern Regional Station
ICAR-National Dairy Research Institute, Adugodi, Bengaluru

ABSTRACT

DPP-IV inhibitors work by blocking the action of DPP-IV, an enzyme which inactivates the incretin hormones. Incretin hormones help the body to produce insulin when it is needed. DPP-IV inhibition is one of the novel approach to treat the diabetes. In this study, focus is given to the milk proteins, as a natural source of DPP-IV inhibition to overcome the side effects of synthetic drugs. Kappa casein was isolated from *Gir* cow milk and analysed for DPP-IV inhibition potential. Enzymatic hydrolysis was done with different enzymes viz., Flavourzyme, pepsin, trypsin and proteinase K for 2-12 hours at different enzyme substrate ratio (1%, 2% and 4%). Trypsin treated kappa casein hydrolysate showed maximum DPP-IV inhibition $83.80 \pm 0.68\%$ after 12 h hydrolysis at 4% E:S ratio. Then hydrolysate with maximum DPP-IV inhibition was passed through ultrafiltration membranes 3 and 10 kDa. Less than 3 kDa fraction showed $73.69 \pm 1.29\%$ DPP-IV inhibition. HPLC analysis was done to collect the fractions from less than 3kDa permeate. Fraction with maximum DPP-IV inhibition was selected for LC-MS/MS analysis to identify the peptides sequence. Two novel peptides with adequate DPP-IV inhibition activity were identified in kappa casein with IC_{50} value 1.79 mM and 1.56 mM. Hence, this hydrolysates rich DPP-IV inhibitors can be a consumed as a dietary supplement, to manage type-2 diabetes.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**STABILITY ENHANCEMENT OF NATURAL
FOOD COLOURS BY MODIFIED PROTEINS**

¹Charu Bisht, ²Neha Rawat and ³S.K. Sharma

Department of Food Science and Technology, COA
GBPUAT, Pantnagar, Uttarakhand

ABSTRACT

Synthetic food colours possess negative health impact due to which consumers are getting attracted towards the use of natural colorants. The four basic natural colours are- anthocyanins, betalains, carotenoids and chlorophylls. Anthocyanins and betalains are water soluble pigments whereas carotenoids and chlorophylls are lipid soluble pigments. Along with imparting colour to the food, these natural pigments show distinct bioactive and therapeutic activities. Lack of stability and less bioavailability of these colorants doesn't promote their utilization even if they are more beneficial. These natural colorants are sensitive towards intrinsic and extrinsic factors. This problem can be tackled by the modifying the proteins. Soya protein fibrillation enhanced the retainment of betalains up to 75% in the food, when processed at 90 °C for 15 minutes. Protein fibrils can be utilized as innovative food material for ameliorating the stability of natural colorants at various temperature, pH, enzymatic activity and level of oxygen. Self-assembled proteins at specific conditions, results in the formation of protein fibrils. These fibril proteins show augmented techno- functional properties. Therefore, they are able to conjugate with the natural colorants. Hence imparting stability to them. Not only the protein fibrils, but also the proteins- polyphenol interaction could stabilize these pigments. The protein interacts with polyphenols either by covalent or non- covalent bonding. Further, these interactions can stabilize the natural colour emulsions. Covalently modified different bovine milk proteins using epigallocatechin- 3- gallate improved the chemical stability of β - carotene in emulsions. Therefore, proteins can be considered as novel ingredient that can help in implementing the utilization of natural colorants.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**METABOLOME PROFILING OF COLOSTRUM, TRANSITION AND
MATURE MILK IN MURRAH BUFFALOES**

**Brijesh, Umesh Kumar, Monika Sodhi, Amarjeet,
Divya Chanda, Abhishek, and Manishi Mukesh**

ICAR-National Bureau of Animal Genetic resources, Karnal-132001

ABSTRACT

Bajaura Makka, H.P. a corn variety grown in Himachal Pradesh with low commercial value was used in this study for recovery of zein and xanthophyll and further application in food products for sustainable development of rural people of HP. The corn meal (CM) and corn gluten meal (CGM) prepared were first hydrolysed using microbial protease, amylase and lipase for extraction of zein at pH 6-10 and optimum pH was 9. The zein content was highest in amylase pre-treated CM (6.47%) and protease pre-treated CGM (4.81%) as detected with HPLC. Highest zein was recovered from both CM (7.14%) and CGM (6.2%) by using mixture of enzymes. Covalent binding was found best immobilization method for protease and lipase at contact time of 3h and 1h, respectively at 30 °C and adsorption method for amylase for 1h at 25 °C. On further treatment highest recovery of zein and xanthophylls was obtained using immobilized protease and amylase in two stage glass bioreactor and further on its filtration through cross flow ultrafiltration system, the relative recovery of zein and xanthophylls increased to 11.81% and 12.35% in CM and 0.10% and 24% in CGM respectively. Presence of different zein fractions was confirmed with SDS-PAGE. Edible zein films were prepared using zein (10% w/w), ethanol (90%), and plasticizers i.e. fructose (4%) and glycerol (5%). The prepared zein films were characterized for physicochemical properties and further used in coating of different model foods viz. Himalayan fig (*Ficus palmata*) leather, roasted nuts, chocolate and vanilla cookies. The extracted xanthophylls was also used as model colorant in yogurt at 0.01 % (w/w) with significant results.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**REACTIVE OXYGEN SPECIES AND DIABETES MELLITUS:
ENVIRONMENTAL IMPACT**

Debidas Ghosh

Professor & HOD, Biomedical Laboratory Science & Management
UGC Research Awardee, Indo-Tunisia Bilateral Fellow
UGC Innovative Funded Dept. Vidyasagar University, Midnapore, West Bengal

ABSTRACT

Environment has a great impact on health maintenance. Environmental factors those deviate the homeostasis of our physiological systems known as stressors and the physiological condition developed in such situation known as stress. Such factors are high temperature and severe cold exposure, different pollutants, food additives etc. Reactive oxygen species (ROS) like singlet O₂, hydrogen peroxide, superoxide anions are formed at high rate in such stressful conditions. As a results pro-oxidant status of our body is dominated over antioxidant status that results health disorder or disease. Diabetes mellitus (DM), a common metabolic cum non-communicable cum life style disease which is now consider as X-syndrome. Environmental stressors can generate free radicals at high level by oxidation-peroxidation chain reaction through ROS generation that results point mutation of gene which is related with production of mutant enzyme, lead to cellular metabolic disorders, especially carbohydrate metabolism, an important cause of diabetes. Moreover, free radicals through cellular signaling system down regulate gene expression of antioxidant enzymes like peroxidase, catalase, glutathione-s-transferase that results high levels of cellular free radicals in liver, kidney and skeletal muscles, most important metabolic organs. As output of this effect, glucose utilization is decreased that lead to DM. Pancreatic β cell degeneration, the source of insulin, is also elevated in such stress full conditions like –unhealthy food style and work style practices that increase the possibility for onset of DM.

For such management of raising level of DM incidence rate throughout the world, natural products having strong antioxidant activities, have major role. It has been noted that nutraceuticals presents in *E.jumbulana*, *Camellia sinensis*, *H. antidysenterica* have major contribution for such management by increasing gene expression of antioxidant enzymes in liver, kidney and skeletal muscles. This field of research unfold a new domain in 21st century about 'Environment and Health' known as 'Neurogenomics'. Moreover, these nutraceuticals also elevate gene expression of hexokinase in liver and kidney that favours glucose utilization in cell. It has also established that such nutraceuticals have ability to generate β cells from hepatic stem cells and so, plasma insulin and c-peptide levels both are elevated. Insulin receptor gene expression is also corrected by such nutraceuticals in diabetic model animals and their efficiency has been comparable with anti-diabetic gold standard drugs. Such studies have been confirmed by genomics and proteomics studies using real time PCR followed by southern blotting and western blotting studies.

It may be concluded that environmental stress factors that results diabetes, a common disease globally and known as 4th killer disease at present, can be managed by natural products of environment, the parallel management process of modern medicine as per guideline of WHO.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



International Conference on
**NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**


30th November - 2nd December, 2022
Goa University, Goa

**SIMULTANEOUS ENANTIOSEPARATION AND SIMULATION STUDIES OF
ATENOLOL, METOPROLOL AND PROPRANOLOL ON CHIRALPAK IG
COLUMN USING SUPERCRITICAL FLUID CHROMATOGRAPHY: A GREEN
APPROACH IN SEPARATION SCIENCE**

Priyanka A. Shah, Pranav S. Shrivastav and Pranav Pandya

Department of Chemistry, School of Sciences
Gujarat University, Navrangpura, Ahmedabad-380009, India

ABSTRACT



Enantioseparation of three β -blockers, atenolol, metoprolol and propranolol is studied on amylose tris(3-chloro-5-methylphenylcarbamate) immobilized chiral stationary phase using supercritical fluid chromatography (SFC). The effect of organic modifiers (methanol, isopropanol and their mixture), column temperature and back pressure on chiral separation of β -blockers was evaluated. Optimum chromatographic separation with respect to resolution, retention, and analysis time was achieved using CO_2 with 25 % methanol: isopropanol (50:50) containing 0.1 % isopropylamine in a single run. Under the optimized conditions, the resolution factors (R_s) and separation factors (α) were greater than 3.0 and 1.5, respectively for all the enantiomers. Further, with increase in temperature (25 to 45 C) and pressure (100 to 150 bars) there was corresponding decrease in retention factors (k), α and R_s . However, a reverse trend (α and R_s) was observed for atenolol with increase in temperature. The thermodynamic data from van't Hoff plots revealed that the enantioseparation was enthalpy driven for metoprolol and propranolol while entropy driven for atenolol. To understand the chiral recognition mechanism and the elution behaviour of the enantiomers, molecular docking studies was performed. The binding energies obtained from simulations studies were in good agreement with the elution order found experimentally and also with the free energy (ΔG) values. The method was validated in the concentration range of 0.5-10 $\mu\text{g/mL}$ for all the enantiomers. The limit of detection and limit of quantitation ranged from 0.126-0.137 $\mu\text{g/mL}$ and 0.376-0.414 $\mu\text{g/mL}$, respectively. The greenness assessment of the method was evaluated by using AGREE software covering all 12 principles of green analytical chemistry. The final score obtained was 0.78, suggesting excellent greenness of the method. Finally, the method was used successfully to analyze these drugs in pharmaceutical preparations.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**STUDIES ON CLINICAL SPECTRUM OF CANINE PYODERMA ASSOCIATED
WITH *STAPHYLOCOCCUS PSEUDINTERMEDIUS* AND OTHER SECONDARY
BACTERIAL INVADERS AND THEIR ANTIBIOGRAM PROFILE FOR
EFFECTIVE THERAPY**

**Prashantha M K; Shambulingappa B E*; Sundareshan S;
Kotresh A M; Rudresh B H; Madhavaprasad C B and Arun S J**

Veterinary College Shimoga-577204 (KVAFSU, Bidar), Karnataka

ABSTRACT

A study on bacteriological investigation of canine pyoderma cases was conducted at the Veterinary College, Shivamogga. Exudate/pus/lesion swabs were collected from clinical cases of canine pyoderma (n=126) and subjected to isolation and identification of bacterial isolates by phenotypic methods. The bacteriological processing of the samples resulted in the recovery of 95 staphylococcal isolates and 18 other bacterial isolates. On culture, staphylococci were the most predominantly (n=95, 75.39%) isolated organisms. The isolates obtained other than staphylococcal species were *E. coli*, *Pseudomonas* species, *Klebsiella* species, *Proteus* species. Subsequently *in vitro* antimicrobial sensitivity testing was carried out for all the isolates. *S. pseudintermedius* showed highest susceptibility to clindamycin and the highest resistance to enrofloxacin, and the other bacterial agents were found sensitive to co-trimoxazole and ceftriaxone and resistant to amoxicillin/sulbactam and enrofloxacin. A study on the occurrence patterns of canine pyoderma revealed higher frequency of cases in male dogs, in the age group of 1-2 years and in Labrador breed. We report the *S. pseudintermedius* as the most predominant pathogen associated with the canine pyoderma in the study area.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**SIMULTANEOUS ESTIMATION OF METHOCARBAMOL AND IBUPROFEN
IN FIXED DOSE COMBINATION BY UV SPECTROPHOTOMETRIC METHOD**

Premanand Velip¹, Anand Mahajan¹, Adison Fernandes¹ and Shweta Borkar^{1*}

¹Department of Pharmaceutical Analysis
Goa College of Pharmacy, Panaj, Goa, India

ABSTRACT

The objective of present research work is to develop a simple, precise and accurate UV spectrophotometric method based on Isobestic point approach for quantitative estimation of methocarbamol (MET) and ibuprofen (IBU) in their fixed dose combinations. All solutions were prepared using methanol as solvent. The isobestic point wavelength for MET and IBU was found to be 226 nm. Concentration of IBU was determined at isobestic wavelength 226 nm while MET was determined directly at 277 nm as there was no interference of IBU. Validation of the method was carried out as per ICH guidelines (Q2R1). Linearity was established for MET at the concentration range of 28 – 56 µg/mL ($r^2 = 0.996$ at 277 nm) and for IBU it was between 16 – 32 µg/mL ($r^2 = 0.990$ at 226 nm). Percentage relative standard deviation (% RSD) value of less than 2 % for intra-day and inter-day precision confirmed the precision of the method. Accuracy of the method was in the range of 98 - 102 %. The developed method is suitable and can be applied for analysis of marketed formulation.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**GREEN SYNTHESIS OF SILVER NANOPARTICLES MEDIATED BY
GARCINIA INDICA (KOKUM) AND THEIR EFFECTIVENESS
AGAINST ACNE-CAUSING BACTERIA**

Trelita de Sousa^{1*} and Gautami S. Teli²

¹Department of Microbiology, St. Xavier's College, Mapusa, Goa – India

²Department of Microbiology, Goa University, Taleigao Plateau, Goa – India

ABSTRACT

Green synthesis of silver nanoparticles (AgNPs) is an emerging field in nanomedicine. *Garcinia indica* (Kokum) is a therapeutic plant indigenous to the Western Ghats of India. In this study, AgNPs were synthesized using aqueous *Garcinia indica* leaf extract as the reducing and capping agent. The AgNPs were characterized by UV-visible absorption spectroscopy, Fourier Transform Infrared (FTIR) spectroscopy, Field Emission Scanning Electron Microscope (FE-SEM) imaging, and Energy dispersive X-ray Diffraction (XRD) analyses. The AgNPs demonstrated both antioxidant and antibacterial properties attributed to the presence of flavonoids and saponins. The antibacterial activity of the AgNPs alone and in combination with ampicillin, chloramphenicol, and rifampicin against acne-causing bacteria revealed a synergistic effect of the AgNPs with all tested antibiotics. Thus, the AgNPs synthesized from *Garcinia indica* leaf extracts using a simple green approach demonstrated promising antioxidant and antibacterial properties which can be further developed for commercial biomedical applications.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**AMELIORATION OF ENVIRONMENTAL TOXICANT BENZO(A) PYRENE
INDUCED TESTICULAR GAMETOGENIC AND STEROIDOGENIC
DYSFUNCTIONS BY A NATURAL ARYL HYDROCARBON RECEPTOR
ANTAGONIST RESVERATROL: INVOLVEMENT OF
P38MAPK/ATF2/INOS SIGNALING**

Kuladip Jana¹, Bhaswati Banerjee¹, Satyajit Halder¹ and Debidas Ghosh²

¹Division of Molecular Medicine, Bose Institute, P1/12, CIT Scheme VIIM

Kolkata-700 054 & EN80, Sector V, Salt Lake City, Bidhannagar, Kolkata-700 091, India

²Department of Biomedical Laboratory Science & Management, Vidyasagar University, Midnapur, India

ABSTRACT

STUDY QUESTION

Does Resveratrol have any protective effect against common environmental toxicant benzo(a)pyrene (B(a)P) induced testicular germ cell apoptosis and dysfunctions of Leydig cell steroidogenesis?

STUDY ANSWER

Resveratrol ameliorates B(a)P induced testicular oxidative stress and germ cell apoptosis by modulating the transcriptional regulation of cytochrome P4501A1 (CYP1A1) and improving Leydig cell steroidogenesis.

BACKGROUND OF THE STUDY

Benzo(a)pyrene, an environmental toxicant mostly present in coal tar, wood burning and car exhaust and which leads to deleterious effects on male reproductive health. Our previous studies have suggested that B(a)P induced oxidative stress, DNA damage and apoptosis involving both mitochondria and death receptor dependent pathways. Natural aryl hydrocarbon receptor (AhR) antagonists may have potential defensive mechanism against B(a)P induced testicular toxicity. The polyphenolic compound resveratrol modulates anti- and pro-apoptotic mediators and document anti-tumorigenic activity. Recently, we have found that resveratrol prevented B(a)P-induced BPDE-DNA adduct formation and spermatogenic dysfunctions in testis. Clinically, resveratrol may be useful to protect against B(a)P induced testicular toxicity, although its effect on the same is not delineated.

STUDY DESIGN, SIZE, DURATION

Forty (40) male Wistar rats (8 weeks; 150–200 gm body weight (b.w.)) were divided into four (n=10); Control (C), Resveratrol (R), Benzo(a)pyrene (B), Benzo(a)pyrene + Resveratrol (B+R) groups those received 0.2 ml corn oil, 50 mg/kg b.w. Resveratrol, 5 mg/kg b.w. Benzo(a)pyrene and 5 mg/kg b.w. Benzo(a)pyrene + 50 mg/kg b.w. Resveratrol respectively by gavages for 60 days. This study was approved by Institutional Animal Ethical Committee (IAEC/BI/08/2016) of Bose Institute, Kolkata, India. (74)

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



International Conference on NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR SUSTAINABLE DEVELOPMENT (NTSD-2022)

30th November - 2nd December, 2022
Goa University, Goa

PARTICIPANTS/MATERIALS, SETTING, METHODS

Testicular germ and Leydig cells were isolated by collagen digestion method. Various molecular biology techniques including immunostaining, Western blot, qRT-PCR, ChIP-assay, flow-cytometry, Terminal-deoxynucleotidyl-transferase-dUTP-nick-end-labeling (TUNEL) and radio-immuno-assay were employed to study the testicular spermatogenesis, steroidogenesis, oxidative stress, apoptosis and B(a)P signaling in order to delineate the ameliorative potential of resveratrol on Benzo(a)Pyrene induced testicular dysfunctions. $P < 0.05$ was considered to be significant as evaluated by One way ANOVA followed by Dunnett multiple comparison test.

MAIN RESULTS AND THE ROLE OF CHANCE

A significant decrease ($P < 0.01$) in epididymal sperm count and motility along with presence of sperm head abnormalities and altered testicular histology were documented following B(a)P treatment. In addition, TUNEL positive apoptotic germ cells were observed with increased ($P < 0.01$) reactive oxygen species generation simultaneously decreasing the activity of antioxidant enzymes and increased mitochondrial damage compared to control. B(a)P disturbed cellular Bax/Bcl2 rheostat, released cytochrome-c, activated caspase 9 and 3 modulating Fas/FasL, Bid and caspase 8 expressions. Steroidogenic-acute-regulatory-protein (StAR) expression and its promoter activation was significantly compromised by B(a)P in Leydig cells. Resveratrol co-treatment with B(a)P maintained testicular redox potential, prevent germ cell apoptosis, increased testosterone level and ameliorated Leydig cell steroidogenic dysfunctions with enhanced expression of CYP1A1, StAR, 3 β HSD and 17 β HSD. Resveratrol also prevented B(a)P induced AhR protein level, its nuclear translocation and subsequent promoter activation of CYP1A1 in germ cells. Resveratrol modulated DAX1 and SF1 expression thus maintaining StAR gene homology and improved Leydig cell steroidogenesis. B(a)P-induced increased iNOS expression through activation of p38MAPK and ATF2, was also ameliorated by Resveratrol. Thus, Resveratrol inhibits conversion of B(a)P to BPDE by modulating the transcriptional regulation of CYP1A1 and acting as antioxidant which prevent germ cell apoptosis and Leydig cell steroidogenesis.

WIDER IMPLICATIONS OF THE FINDINGS

Resveratrol being an antioxidant and a natural AhR antagonist, works as a double-edged sword against environmental toxicant B(a)P induced male reproductive toxicity. Present day B(a)P induced toxicokinetics and its protection is not well elucidated in human. Resveratrol can come up with new hopes as potential protector of reproductive health against B(a)P.

STUDY FUNDING

This study was supported by the award of DST First Track Young Scientist Scheme of Govt. of India to Dr. Kuladip Jana.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**A NOVEL CALEBIN A ANALOG INDUCES ROS AND INHIBITS
CANCER PROGRESSION VIA DISRUPTING PI3K/AKT PATHWAY**

Satyajit Halder¹, Pradip Shit¹, Ananya Dutta^{1,2}, Anup Kumar Misra^{1*} and Kuladip Jana^{1*}

¹Division of Molecular Medicine, Bose Institute, P-1/12, Calcutta Improvement Trust Scheme VII M, Kolkata-700054 and EN 80, Sector V, Salt Lake City, Bidhannagar, Kolkata-700091, India.

²Department of Pathology and Laboratory Medicine, The University of North Carolina, Chapel Hill, U.S.A.

ABSTRACT

Chemotherapy, irradiation and immunotherapy emerges as the gold standard approaches for the treatment of cancer worldwide but due to their toxicity they have some limitations. Thus due to the toxicity of currently used therapeutics for the treatment of various types of tumors, use of natural products from plants and animals and marine organisms and their derivatives for proper targeting of both the cancer cell and cancer stem cell colonies have produced remarkable leads for the control of cancer due to their limited cytotoxicity to the normal cell as compared to cancer cells. Natural yields and their structural derivatives (small molecules) have contributed to nearly 50% of the drugs used for cancer chemoprevention. Calebin-A, a naturally occurring curcuminoid analog derived from turmeric root (*Curcuma longa*) has gained attention due to its tremendous anti-cancer potential. Thus, keeping this in mind an array of Calebin-A analogs were designed and created and were evaluated for their cytotoxic potentials on various murine and human malignant cell lines. One such analog compound AS-18 was effectively opted out as most potent with a high cytotoxic effect against cancer cells and less toxicity against normal cell lines. Our present study established ROS-induced mitochondrial dysfunction and finally apoptosis induction by the lead compound. The inhibition of cell propagation was linked to the data confirming G2/M phase arrest. The mitochondrial dysfunction was confirmed by JC1 and MitoSOX assay along with the change in Bax/Bcl-2 ratio. The compound also blocked the PI3K/Akt pathway. Molecular docking and western blot investigation established these pieces of information. The lead compound AS-18 suppressed the NRF-2 protein expression thus increasing the free radicals in the tumor cells. The compound AS-18 induced ROS-mediated caspase-dependent apoptosis as the western blot data confirmed caspase activation. The 4T1 injected Balb/C syngeneic tumor model confirmed the augmentation in the inhibitory outcome of the lead compound. This study sums up the mechanistic pathway by which the compound AS-18 mediates its cytotoxic effect in cancer cells. A novel Calebin-A compound AS-18 acted as an antagonist against the IGF R1 receptor, blocked the PI3K/Akt and NRF-2 pathway, and encouraged apoptosis via ROS generation.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**EVALUATION OF METAL TOLERANCE OF HALOPHILIC
EUBACTERIAL AND ARCHAEAL STRAINS ISOLATED
FROM SOLAR SALTERNS OF GOA INDIA**

Fazila Shaikh^{1*} and Sanika Naik-Samant¹

¹Biotechnology Discipline
School of Biological Sciences & Biotechnology, Goa University

ABSTRACT

Marine econiche is continuously exposed to metal pollution thus enabling estuaries and solar salterns to serve as an effective sink for these metals, leading to their high accumulation through evaporation. Estuarine environments thus serve as a good source of metal tolerant microbes which also contribute towards the biogeochemical cycling of inorganic elements. In this study, the halophilic strains from solar salterns of Agarvado and Maina, Goa-India were obtained on NTYE and ZMA agar plates incorporated with 25 % crude salt. The halophilic isolates were characterized based on their Gram character; tolerance to salt, antibiotic sensitivity, response to sodium taurocholate and evaluation of the pigment extracted. The potential eubacterial and haloarchaeal strains were screened for their metal tolerance/resistance ability using metal salts of Cu, Ni and Ag at 0.5, 1, 1.5, 2 and 4 mM concentrations. Tolerance of halophilic isolates to metal salts was observed by monitoring the growth for each of the culture on NTYE plates amended with the respective metal salts for an incubation period of 5-7 days. Growth and effect on pigment production of halophilic isolates in NTYE medium in absence and presence of 1 mM concentration of different metal ions was also analyzed. Culture tolerating/resisting metal salt exhibited alteration in pigment production. This indicated the role of cell wall components of potential cultures in withstanding the metal ions. Thus, showing metal tolerance in halophilic isolates isolated from solar salterns. Further, the potential cultures can also be used in synthesis of metal nanoparticles of nanobiotechnological significance along with the bioremediation applicability in metal contaminated saline soils.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by



**International Conference on
NATURAL SCIENCE AND GREEN TECHNOLOGIES FOR
SUSTAINABLE DEVELOPMENT (NTSD-2022)**

**30th November - 2nd December, 2022
Goa University, Goa**

**PERFORMANCE OF PLANT SPECIES ON
THE TEXTILE EFFLUENT AFFECTED SOIL**

**Dipak Kumar Gupta^{1*}, Anil Kumar Shukla²,
Kamla K Choudhary², SR Meena² and Keerthika A.**

¹ICAR-India Agricultural Research Institute-Jharkhand, Gauria Karma, Hazaribagh

²ICAR-Central Arid Zone Research Institute, RRS, Pali-Marwar, Rajasthan

ABSTRACT

Pali district of Rajasthan is one of the textile (printing and dyeing) industrial hubs of India. Textile industries are high water requiring units and consume about 200 litres of water for processing of 1 kg of textile and thus this industry generate large amount of effluent. The treated and untreated effluent from textile industries in Pali, was directly or indirectly discharged into Bandi River. Agricultural land downstream of Bandi River from Pali city has been reported to be degraded due to irrigation from textile effluent contaminated Bandi River. These agricultural land were ones used for the cultivation of crops like wheat, mustard, vegetables etc. is presently not been cultivated from last 25-30 years. With this background, present study evaluated the growth of seven plant species on this contaminated soil at CAZRI-Regional Research Station, Pali, Marwar, with aim to re-cultivate the land. *Parkinsonia aculeate* (Parkinsonia), *Ailanthus excelsa* (Ardu), *Lawsonia inermis* (Henna), *Suaeda fruticosa* (Suaeda), *Cymbopogon martini* (Palmarosa), *Vetiveria zizanioides* (Vetiver) and *Diplachne fusca* (Karnal grass) were planted in pots filled with affected soil and amended with various amendments. After three month of planting, the plant growth parameters were recorded. The result showed significant difference in growth parameters among plant species. Out of seven plant species evaluated on textile effluent affected soil, *Diplachne fusca* (Karnal grass), *Vetiveria zizanioides* (Vetiver aromatic plant), *Lawsonia inermis* (Henna) and *Cymbopogon martini* (Palmarosa) performed well on textile effluent affected soil in terms of plant growth and biomass yield in both the years (2018 and 2019) of experiment.

Jointly Organized by



National Environmental
Science Academy, New Delhi



School of Biological Sciences and Biotechnology
Goa University, Taleigao Plateau, Goa

Sponsored by





INVITATION OF RESEARCH ARTICLES for PUBLICATION in NESA Journals

INTERNATIONAL JOURNAL ON AGRICULTURAL SCIENCES

ISSN NO. 0976-450X | NAAS RATING 2.60

INTERNATIONAL JOURNAL ON ENVIRONMENTAL SCIENCES

ISSN NO. 0976-4534

INTERNATIONAL JOURNAL ON BIOLOGICAL SCIENCES

ISSN NO. 0976-4518

INDIAN JOURNAL OF UNANI MEDICINE

ISSN NO. 0974-6056

These JOURNALS ON DIFFERENT SUBJECTS are being published by this Academy. Send your manuscripts for peer-review by online / e-mail. THE AUTHORS MUST MENTION ADDRESS, Contact Nos. and E-MAIL ID in their forwarding letter. Proof will be sent for correction before publishing. A pledge for originality will be signed by the authors. A soft copy of final manuscript will be sent by mail and journal copy by post after the receipt of the **PROCESSING FEE**. All remittances can be transfer through NEFT/RTGS in below bank detail:

Name of the Bank: Bank of Maharashtra, Kalkaji Branch, New Delhi – 110 019 (INDIA)

Name of Account Holder: NATIONAL ENVIRONMENTAL SCIENCE ACADEMY

Account Number: 20066872035:

IFSC Code: MAHB0000974

in favour of **NATIONAL ENVIRONMENTAL SCIENCE ACADEMY (NESA), NEW DELHI.**

For further details and **NOTES FOR AUTHORS**,
please contact Academy at
nesapublications@gmail.com infonesa88@gmail.com



NATIONAL ENVIRONMENTAL SCIENCE ACADEMY

206, Raj Tower-1, Alaknanda Community Centre, New Delhi - 110 019

Tel.: 011-2602 3614 • M: 9811238475, 9811238995, 9971383650

E-mail: infonesa88@gmail.com; neaspublishations@gmail.com

Website: www.nesa-india.org

